

# Phytochemical Constituents and Pharmacological Activities of *Eryngium* L. (Apiaceae)

Ping Wang, Zushang Su, Wei Yuan, Guangrui Deng and Shiyu Li\*

National Center for Pharmaceutical Crops, Arthur Temple College of Forestry and Agriculture, Stephen F. Austin State University, Nacogdoches, TX, 75962-6109, USA

**Abstract:** *Eryngium* L. is the largest and arguably the most taxonomically complex genus of the family Apiaceae. The genus has approximately 250 species throughout the world, with the center of diversity in South America. Some *Eryngium* species are cultivated as ornamental, vegetable, or medicinal crops for folk uses. With increasing chemical and biological investigations, *Eryngium* has shown its potential as pharmaceutical crops. This review focuses on phytochemistry and pharmacological activities of 127 compounds isolated and identified from 23 species of *Eryngium*, particularly non-essential oil compounds such as terpenoids, triterpenoid saponins, flavonoids, coumarins, polyacetylenes, and steroids. *Eryngium* extracts or isolates have shown *in vitro* bioactivities such as cytotoxicity against various human tumor cell lines, anti-inflammatory, anti-snake and scorpion venoms, antibacterial, antifungal, and antimalarial, antioxidant, and antihyperglycemic effects. *In vivo* studies through various animal models have also shown promising results. However, chemical constituents and their bioactivities of most species of this highly diversified genus have not been investigated. The molecular mechanism of bioactivities (particularly cytotoxicity and anti-snake and scorpion venoms) of *Eryngium* isolates remains elusive. Also, anti-tumor activity of polyhydroxylated triterpenoid saponins isolated from *Eryngium* needed to be further explored.

**Keywords:** *Eryngium* L., ethnobotany, phytochemistry, triterpenoid saponins, flavonoids, polyacetylenes, cytotoxicity, anti-inflammatory activity, anti-snake and scorpion venoms effects.

## INTRODUCTION

*Eryngium* L. consists of approximately 250 species in Eurasia, North Africa, North and South America, and Australia [1, 2]. It is the largest and arguably the most taxonomically complex genus of the family Apiaceae [3]. Some species are rare or endangered, e.g., *E. alpinum* L., *E. aristulatum* Jeps. var. *parishii* (J.M. Coult. & Rose) Mathias & Constance, *E. constancei* Sheikh, *E. cuneifolium* Small, *E. viviparum* [4-7]. Wolff's (1913) treatment of *Eryngium* is the most comprehensive and predominant: two groups were recognized within the genus *Eryngium*: "Species gerontogaeae" representing 12 sections from the Old World (Africa, Europe, and Asia) and "Species americanae and australienses" including 22 sections from the New World (Americas and Australia) [8]. Based on morphology, Wörz (2005) proposed five subgenera within *Eryngium*: subg. *Eryngium*, subg. *Fruticosa*, subg. *Monocotyloidea*, subg. *Semiaquatica*, and subg. *Foetida* with the subg. *Eryngium* occurring in Europe, Western Asia and North Africa, the other four subgenera mostly in the New World and in Australia [2, 9]. However, Wörz's classification is not supported by phylogenetic data. Recent infrageneric relationship analysis of Calviño *et al.* (2007) using sequence data from the chloroplast DNA trnQ-trnK 5'-exon and

nuclear ribosomal DNA ITS regions of 118 species support Wolff's two-group classification [3]. Calviño *et al.* (2007) recognized two subgenera within the genus: subg. *Eryngium* and subg. *Monocotyloidea* [3].

Like many other members of the celery or carrot family, *Eryngium* has been used as ornamental, vegetable, or medicinal plants. Some species, such as *E. foetidum* L., *E. maritimum* L., *E. planum* L., *E. dichotomum* Desf., *E. campestre* L. and *E. creticum* Lam. have been used as food or in traditional medicine locally or worldwide [10]. *E. foetidum* and *E. caucasicum* Trautv. are cultivated as leaf vegetable crops in Asia and Africa [11, 12]. The fruits of *E. foetidum* were taken as food in Nigeria. The plant is indigenous to Tropical America and the West Indies where it is used as medicine and food some having domesticated the plant in their kitchen gardens and orchards. It has become naturalized and often is cultivated across South Asia, the Pacific islands, Tropical Africa and the warmer southern parts of Europe [13].

Some species have been used in folk medicine. *E. campestre* is a well-known plant of the Apiaceae family and is used in Turkish folk medicine. Infusions of the aerial and root parts are used as an antitussive, diuretic, appetizer, stimulant, and aphrodisiac [14-16]. *E. creticum* has been used in folk medicine in Jordan as a remedy for scorpion stings in the rural areas and as a hypoglycemic agent [17]. *E. elegans* Cham. & Schltdl. was reported to be used for diuretic uses in Argentina [18], and *E. foetidum* for the treatment of several anti-inflammatory disorders in China [19].

\*Address correspondence to this author at the National Center for Pharmaceutical Crops, Arthur Temple College of Forestry and Agriculture, Stephen F. Austin State University, Nacogdoches, TX 75962, USA; Tel: 936-468-2071; Fax: 936-468-7058; E-mail: lis@sfasu.edu

Several *Eryngium* species are used as medicine by various tribes of Native Americans. Whole plants of *E. alismifolium* are used for diarrhea; roots of *E. aquaticum* are used as emetic and gastrointestinal infusion, antidote for poisons, tapeworms and pinworms, diuretic, and venereal diseases; and roots of *E. yuccifolium* Michx. are used as snakebite and toothache remedy as well as for neuralgia, bladder and kidney troubles; roots of *E. yuccifolium* var. *synchaetum* are used for human and animal sickness such as digestive problems, diarrhea, headache, body soreness, and snakebites [20]. Some recent bioactivity investigations have confirmed some traditional medicinal uses. Ethanol extracts of *E. billardieri* Delar., *E. campestre*, *E. creticum*, *E. davisii*, *E. foetidum*, *E. isauricum*, *E. kotschyi*, *E. maritimum*, and *E. trisectum* showed apparent anti-inflammatory and antinociceptive activity [21-23]. The fresh leaf extract of *E. creticum* gave a higher percentage inhibition of the haemolytic activity of the scorpion venom compared with the dried leaf extract, but extracts of both fresh and dried roots of *E. creticum* gave 100% inhibition of the snake and scorpion venoms [24]. Also, some species (i.e., *E. caucasicum*) showed antioxidant activity [25, 26].

To date, terpenoids, triterpenoid saponins, flavonoids, coumarins, polyacetylenes, steroids, and essential oils have been reported in the genus *Eryngium*. However, most species of *Eryngium* have not been extensively investigated in chemical constituents. Existing phytochemical investigations indicated the presence of essential oils (primarily sesquiterpenes and monoterpenes) in *E. bourgatii* Gouan (*E. amethystinum* Lam.), *E. billardieri*, *E. bourgatii* Gouan, *E. bungei* Boiss., *E. caeruleum* M. Bieb., *E. campestre*, *E. corniculatum*, *E. creticum*, *E. foetidum*, *E. giganteum* M. Bieb., *E. glaciale* Boiss., *E. palmatum* Vis. et Pančić, *E. paludosum* (Moore et Betche) Michael, *E. paniculatum* Cav. & Domb. ex Delar., *E. rosulatum* Michael, *E. serbicum* Pančić, *E. vesiculosum* Labill., and *E. yuccifolium* [25, 27-48]. Other classes of compounds isolated from *Eryngium* include flavonoids from *E. campestre*, *E. giganteum*, *E. macrocalyx* Schrenk, *E. maritimum*, *E. octophyllum* Eug. Kor., and *E. yuccifolium* [10, 49-54], coumarins from *E. campestre* and *E. ilicifolium* Lam. [55-57], sterols from *E. foetidum* and *E. agavifolium* [21], a rosmarinic acid derivative from *E. alpinum* L. [13], lactone from *E. carlinae* Delar. [58], and triterpenoid glycosides from *E. bourgatii*, *E. bromeliifolium* Delar., *E. campestre*, *E. giganteum*, *E. macrocalyx*, *E. maritimum*, *E. octophyllum*, *E. planum*, and *E. yuccifolium* [10, 14, 15, 59-66]. The main saponins from this genus belong to polyhydroxylated triterpenoid glycosides with ester functions. This class of saponin has been found in various groups of plants such as *Aesculus chinensis* L. (Hippocastanaceae) [67-71], *Pittosporum tobira* (Thunb.) Ait. (Pittosporaceae) [72], *Sanicula elata* var. *chinensis* Makino (Apiaceae) [73], and *Harpullia austro-caledonica* Baill. (Sapindaceae) [74]. This type of saponin has been shown to possess anti-inflammatory property [71, 75, 76], anti-HIV-1 protease activity [68], and cytotoxicity for tumor cells [72, 77-79]. Recently, we isolated and identified 25 new polyhydroxylated triterpenoid saponins from North American *Aesculus pavia* L. [77, 80]. The saponins with two acyl groups at C-21 and C-22 had cytotoxic activity against 60 cell lines from nine different human cancers [77, 81].

This review focuses on phytochemistry and pharmacological activities of 127 compounds isolated and identified from *Eryngium*, particularly non-essential oil compounds such as terpenoids, triterpenoid saponins, flavonoids, coumarins, polyacetylenes, and steroids. This involves 23 species belonging to 9-11 sections of *Eryngium* in both Old and New Worlds (Table 1).

## PHYTOCHEMISTRY

Of 250 species of the genus *Eryngium*, only 23 species have been more or less investigated phytochemically. To date, at least 127 compounds, primarily phenolic compounds and terpenoids have been isolated and identified from these species, including triterpenoid saponins, monoterpene, sesquiterpenes, triterpenoids, flavonoids, coumarins, steroids, acetylenes, and other classes of compounds. A summary of the compounds isolated from *Eryngium* species is carried in Tables 2 and 3.

### Triterpenoid Glycosides

The genus *Eryngium* is known to contain triterpenoid saponins as the main components. To date, 25 saponins have been isolated from this genus. The most of *Eryngium* saponins belong to polyhydroxylated oleanene triterpenoid saponins (Table 2). This class of saponins has been found in a wide range of plants belonging to different families, such as *Aesculus* L. (Hippocastanaceae) [100], *Pittosporum tobira* (Thunb.) Ait. (Pittosporaceae), and *Harpullia austro-caledonica* Baill. (Sapindaceae) [10]. In the family Apiaceae, *Hydrocotyle* L., *Hacquetia* Neck ex DC., *Steganotaenia* Hochst, and *Sanicula* L. have already indicated the present of polyhydroxylated triterpenoid saponins [14].

Recently, three detailed phytochemical investigation on the roots of *E. campestre* [14, 15] and the whole plant of *E. yuccifolium* [10] resulted in the isolation and structural elucidation of 19 new polyhydroxylated oleanene triterpenoid saponins, named 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22-*O*- $\beta$ , $\beta$ -dimethylacryloyl-A<sub>1</sub>-barrigenol (1), 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22-*O*-angeloyl-R<sub>1</sub>-barrigenol (2), 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-21-*O*-acetyl-22-*O*-angeloyl-R<sub>1</sub>-barrigenol (3), 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-21-*O*-acetyl-22-*O*- $\beta$ , $\beta$ -dimethylacryloyl-R<sub>1</sub>-barrigenol (4), 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22-*O*-angeloyl-28-*O*-acetyl-R<sub>1</sub>-barrigenol (5), 3-*O*- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-[ $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)]- $\beta$ -D-glucuronopyranosyl-22-*O*-angeloyl-R<sub>1</sub>-barrigenol (6), 3-*O*- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-[ $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)]- $\beta$ -D-glucuronopyranosyl-22-*O*- $\beta$ , $\beta$ -dimethylacryloyl-A<sub>1</sub>-barrigenol (7), eryngiosides A-L (8-19), together with two known saponins 21 $\beta$ -angeloyloxy-3 $\beta$ -[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]-[ $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 3)]- $\beta$ -D-glucuronopyranosyloxyolean-12-ene-15 $\alpha$ , 16 $\alpha$ , 22 $\alpha$ , 28-tetrol (20) and saniculasaponin III (21).

Four triterpenoids R<sub>1</sub>-barrigenol (28), A<sub>1</sub>-barrigenol (29), barringtogenol C (31), and 3 $\beta$ , 16 $\alpha$ , 22 $\alpha$ , 28-tetrahydroxyolean-12-ene derivatives were recognized to be the aglycones of polyhydroxylated *Eryngium* saponins with acyl functions at C-21 and C-22 positions. The *Eryngium* saponins resemble that of *Aesculus* species, but there were

Table 1. List of *Eryngium* Species Included in this Review.

Section	Species	Reference
<b>OLD WORLD</b>		
Section Alpina	<i>E. alpinum</i> L.	[13]
	<i>E. giganteum</i> M. Bieb. (synonyms: <i>E. asperifolium</i> F. Delar., <i>E. glaucum</i> Hoffm.)	[13, 46, 54]
Section Campestris	<i>E. bourgatii</i> Gouan (synonyms: <i>E. alpinum</i> Lapeyr., <i>E. amethystinum</i> Lam., <i>E. pallens</i> Mill., <i>E. planum</i> Lapeyr., <i>E. tounefortii</i> Bubani)	[13, 46, 82-84]
	<i>E. campestre</i> L. (synonyms: <i>E. latifolium</i> Hoffm. & Link ex Willk & Lange, <i>E. officinale</i> Garsault, <i>E. trifidum</i> L.)	[14, 15, 45, 50, 52, 55, 85]
Section Dryophylla	<i>E. ilicifolium</i> Lam.	[56]
	<i>E. octophyllum</i> Korovin (synonyms: <i>E. mirandum</i> Bobrov)	[51, 66]
Section Halobia	<i>E. macrocalyx</i> Schrenk (synonym: <i>E. incognitum</i> Paviov)	[65, 86, 87]
	<i>E. maritimum</i> L. (synonym: <i>E. marinum</i> Garsault)	[46, 49, 63, 88]
Section Palmito	<i>E. serbicum</i> Pančić	[82]
Section Plana	<i>E. caucasicum</i> Trautv. (synonyms: <i>E. biebersteinianum</i> Nevski)	[66, 89]
	<i>E. creticum</i> Lam. (synonyms: <i>E. cyaneum</i> Sm., <i>E. syriacum</i> Lam.)	[25, 90, 91]
	<i>E. dichotomum</i> Desf.	[53, 92]
	<i>E. planum</i> L. (synonyms: <i>E. E. armatum</i> Csátó ex Simonk., <i>E. caeruleum</i> M. Bieb., <i>E. dalla-torrei</i> M. Hiroe, <i>E. intermedium</i> Weinm.), <i>E. latifolium</i> Gilib., <i>E. planifolium</i> Pall., <i>E. planum</i> Lindl., <i>E. planum</i> var. <i>armatum</i> Csátó ex Simonk., <i>E. pumilum</i> Gilib., <i>E. pusillum</i> Gilib.)	[13, 46, 86, 93-98]
	<i>E. variifolium</i> Coss.	[13, 46]
	<i>E. spinalba</i> L. (synonyms: <i>E. rigidum</i> Lam., <i>E. leucacanthum</i> St.-Lag.)	[13]
Section (unknown)	<i>E. tripartitum</i> Desf.	[13]
<b>NEW WORLD</b>		
Section Areata	<i>E. agavifolium</i> Griseb.	[13]
Section Foetida	<i>E. foetidum</i> L. (synonyms: <i>E. antihystericum</i> Rottler, <i>E. antihystericum</i> Rottb.)	[21, 59, 99]
Section (unknown)	<i>E. bromeliifolium</i> Delar.	[60-62]
Section Panniculata	<i>E. eburneum</i> Decne. (synonyms: <i>E. bracteosum</i> (DC.) Griseb., <i>E. paniculatum</i> var. <i>bracteosum</i> DC.)	[13]
	<i>E. pandanifolium</i> Cham. & Schldl. (synonyms: <i>E. decaisneanum</i> Urb., <i>E. oligodon</i> (DC.) Griseb., <i>E. pandanifolium</i> var. <i>atrocephalum</i> Kuntze), <i>E. paniculatum</i> var. <i>oligodon</i> DC.)	[13]
	<i>E. paniculatum</i> Cav. & Domb. Ex Delar. (synonyms: <i>E. paniculatum</i> var. <i>chinense</i> DC., <i>E. paniculatum</i> f. <i>junior</i> Urb., <i>E. paniculatum</i> var. <i>litorale</i> G. Kunkel, <i>E. subulatum</i> Vell.)	[47]
	<i>E. yuccifolium</i> Michx.	[10, 13, 27]

various differences between both saponins in aglycons and sugar components. Especially, all saponins identified from *E. campestre* are interesting structurally because the acylation by a  $\beta,\beta$ -dimethylacrylic acid or angeloyl at the C-22 position is rare among triterpenoid saponins [14]. *Eryngium* saponins from *E. yuccifolium* are characterized as an angeloyl (Ang) attached to C-21 or C-22, and an angeloyl (Ang)/acetyl (Ac) group to the C-21/22 positions. The saponins structurally based on three aglycones of R<sub>1</sub>-barrigenol (**28**), A1-barrigenol (**29**), barringtogenol C (**31**) with the hydroxyl group at C-15 and/or C-21 contained one oligosaccharide chain with a glucuronopyranosyl unit attached to C-3 position of the aglycones. This oligosaccharide chain had a rhamnopyranosyl or an arabinopyranosyl unit attached to the glucuronopyranosyl unit. The saponins pos-

sessed the aglycone of 3 $\beta$ , 16 $\alpha$ , 22 $\alpha$ , 28-tetrahydroxyolean-12-ene without a hydroxyl group at C-15 and C-21 linked a trisaccharide chain of glucoses and galactoses to the C-3 position. However, these saponins were identified bearing none of acyl functions at C-21 and C-22. It was believed that at least one angeloyl moiety at either C-21 or C-22 positions of *Eryngium* saponins by a recent structure-activity relationship (SAR) investigation makes significant contribution to the cytotoxicities, but the type and number of sugar moiety at C-3 may decrease their cytotoxicities [101].

Some other classes of triterpenoid saponins were also reported in *Eryngium* species. 3-O- $\beta$ -D-glucopyranosyl oleanolic acid 28-O- $\beta$ -D-xylopyranoside (**22**) and 3-O- $[\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)- $[\beta$ -D-fucopyranosyl-(1 $\rightarrow$ 3)]- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-glucopyranosyl]-olean-12-en-

**Table 2. Major Non-Essential oil Compounds Isolated and Identified from *Eryngium***

No.	Compound Name	Botanical Source	Reference
	<b>Triterpenoid Saponins</b>		
1	3- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22- <i>O</i> - $\beta$ , $\beta$ -dimethylacryloyl-A <sub>1</sub> -barrigenol	<i>E. campestre</i>	[15]
2	3- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22- <i>O</i> -angeloyl-R <sub>1</sub> -barrigenol	<i>E. campestre</i>	[15]
3	3- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-21- <i>O</i> -acetyl-22- <i>O</i> -angeloyl-R <sub>1</sub> -barrigenol	<i>E. campestre</i>	[15]
4	3- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-21- <i>O</i> -acetyl-22- <i>O</i> - $\beta$ , $\beta$ -dimethylacryloyl-R <sub>1</sub> -barrigenol	<i>E. campestre</i>	[15]
5	3- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucuronopyranosyl-22- <i>O</i> -angeloyl-28- <i>O</i> -acetyl-R <sub>1</sub> -barrigenol	<i>E. campestre</i>	[15]
6	3- <i>O</i> - $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-[ $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)]- $\beta$ -D-glucuronopyranosyl-22- <i>O</i> -angeloyl-R <sub>1</sub> -barrigenol	<i>E. campestre</i>	[14]
7	3- <i>O</i> - $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-[ $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)]- $\beta$ -D-glucuronopyranosyl-22- <i>O</i> - $\beta$ , $\beta$ -dimethylacryloyl-A <sub>1</sub> -barrigenol	<i>E. campestre</i>	[14]
8	eryngiosides A	<i>E. yuccifolium</i>	[10]
9	eryngiosides B	<i>E. yuccifolium</i>	[10]
10	eryngiosides C	<i>E. yuccifolium</i>	[10]
11	eryngiosides D	<i>E. yuccifolium</i>	[10]
12	eryngiosides E	<i>E. yuccifolium</i>	[10]
13	eryngiosides F	<i>E. yuccifolium</i>	[10]
14	eryngiosides G	<i>E. yuccifolium</i>	[10]
15	eryngiosides H	<i>E. yuccifolium</i>	[10]
16	eryngiosides I	<i>E. yuccifolium</i>	[10]
17	eryngiosides J	<i>E. yuccifolium</i>	[10]
18	eryngiosides K	<i>E. yuccifolium</i>	[10]
19	eryngiosides L	<i>E. yuccifolium</i>	[10]
20	21 $\beta$ -angeloyloxy-3 $\beta$ -[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]-[ $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 3)]- $\beta$ -D-glucuronopyranosyloxyolean-12-ene-15 $\alpha$ , 16 $\alpha$ , 22 $\alpha$ , 28-tetrol	<i>E. yuccifolium</i>	[10]
21	saniculasaponin III	<i>E. yuccifolium</i>	[10]
22	3- <i>O</i> - $\beta$ -D-glucopyranosyl oleanolic acid 28- <i>O</i> - $\beta$ -D-xylopyranoside	<i>E. bromeliifolium</i>	[60]
23	3- <i>O</i> -[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)]-[ $\beta$ -D-fucopyranosyl-(1 $\rightarrow$ 3)]- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-glucopyranosyl]-olean-12-en-23,28-diol	<i>E. foetidum</i>	[59]
24	betulinic acid 3- <i>O</i> - $\beta$ -D-glucopyranoside	<i>E. bromeliifolium</i>	[62]
25	betulinic acid-3- <i>O</i> - $\beta$ -D-glucopyranosyl(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside	<i>E. bromeliifolium</i>	[61]
	<b>Triterpenoids</b>		
26	betulinic acid	<i>E. bromeliifolium</i>	[64]
27	oleanolic acid	<i>E. macrocalyx (E. incognitum)</i>	[65, 87]

Table 2. Contd....

No.	Compound Name	Botanical Source	Reference
28	R <sub>1</sub> -barringenol	<i>E. maritimum</i> <i>E. planum</i> <i>E. caucasicum</i> ( <i>E. biebersteini-anum</i> ) <i>E. octophyllum</i>	[63, 66, 95]
29	A <sub>1</sub> -barringenol	<i>E. maritimum</i> <i>E. planum</i> <i>E. caucasicum</i> ( <i>E. biebersteini-anum</i> ) <i>E. octophyllum</i>	[63, 66, 95]
30	22-(2-methyl-2-butenolate)-olean-12-ene-3,15,16,22,28-pentol	<i>E. macrocalyx</i>	[65]
31	barringtogenol C	<i>E. maritimum</i> <i>E. planum</i>	[63, 95]
32	erynginol A	<i>E. planum</i>	[95]
33	22 $\alpha$ -hydroxyerythrodiol	<i>E. maritimum</i>	[63]
	<b>Sesquiterpenes</b>		
34	eryng-9-en-15-al	<i>E. creticum</i>	[25]
35	15-hydroxy- $\alpha$ -muurolene	<i>E. giganteum</i>	[46]
36	15-oxy- $\alpha$ -muurolene	<i>E. giganteum</i>	[46]
37	15-nor- $\alpha$ -muurolene	<i>E. giganteum</i>	[46]
38	(+)-ledol	<i>E. giganteum</i>	[46]
39	(+)-spathulenol	<i>E. giganteum</i>	[46]
40	germacrene D	<i>E. giganteum</i>	[46]
41	trans- $\beta$ -farnesene	<i>E. giganteum</i>	[46]
	<b>Monoterpenes</b>		
42	3-( $\beta$ -D-glucopyranosyloxymethyl)-2,4,4-trimethyl-2,5-cyclohexadien-1-one	<i>E. campestre</i> <i>E. creticum</i>	[45, 91]
43	3-( $\beta$ -D-glucopyranosyloxymethyl)-2,4,4-trimethyl-2-cyclohexen-1-one	<i>E. campestre</i>	[45]
44	5-[( $\beta$ -D-glucopyranosyloxy)methyl]-4-hydroxy-4-[(1 <i>E</i> ,3 <i>S</i> )-3-hydroxy-1-butenyl]-3,5-dimethyl-2-cyclohexen-1-one,	<i>E. dichotomum</i>	[53]
45	isoferulyl senecioate	<i>E. variifolium</i>	[46]
46	(-)-2,4,4-trimethyl-3-formyl-2,5-cyclohexadienyl angelate	<i>E. paniculatum</i>	[47]
47	<i>O</i> -[2-angeloyloxymethyl- <i>cis</i> -crotonoyl]-ferulol	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
48	<i>O</i> -[2-angeloyloxymethyl- <i>cis</i> -crotonoyl]-isoferulol	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
49	<i>O</i> -[2-(2-methyl-butyryloxymethyl)- <i>cis</i> -crotonoyl]-ferulol	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
50	<i>O</i> -[2-(2-methyl-butyryloxymethyl)- <i>cis</i> -crotonoyl]-isoferulol	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]

Table 2. Contd....

No.	Compound Name	Botanical Source	Reference
51	<i>O</i> -[2-isovaleryloxymethyl)- <i>cis</i> -crotonoyl]-isoferulol	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
	<b>Flavonoids</b>		
52	quercetin	<i>E. creticum</i>	[91]
53	quercitrin	<i>E. campestre</i>	[52]
54	isoquercitrin	<i>E. campestre</i> <i>E. maritimum</i> <i>E. giganteum</i>	[49, 52, 54]
55	rutin	<i>E. campestre</i> <i>E. octophyllum</i>	[51, 52]
56	luteolin 7- <i>O</i> - $\beta$ -D-glucopyranoside	<i>E. campestre</i>	[52]
57	astragalin	<i>E. campestre</i> <i>E. maritimum</i>	[49, 52, 54]
58	kaempferol 7- <i>O</i> - $\alpha$ -L-rhamnopyranoside	<i>E. campestre</i>	[52]
59	kaempferol 3- $\beta$ -D-glucopyranosyl-7- <i>O</i> - $\alpha$ -L-rhamnopyranoside	<i>E. maritimum</i> <i>E. macrocalyx</i>	[49, 51]
60	kaempferol 3,7-di- <i>O</i> - $\alpha$ -L-rhamnopyranoside	<i>E. campestre</i> <i>E. planum</i> <i>E. giganteum</i>	[52, 54, 97, 98]
61	kaempferol-3- <i>O</i> -(6- <i>O</i> - $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranoside	<i>E. planum</i>	[96]
62	kaempferol 3- <i>O</i> - $\beta$ -D-(2'- <i>p</i> - <i>E</i> -hydroxycinnamoyl) -glucopyranoside	<i>E. campestre</i>	[50]
63	kaempferol 3- <i>O</i> - $\beta$ -D-(2'- <i>p</i> - <i>Z</i> -hydroxycinnamoyl)- glucopyranoside	<i>E. campestre</i>	[50]
64	kaempferol-3- <i>O</i> -(2- <i>O</i> - <i>trans</i> - <i>p</i> -methoxycoumaroyl-6- <i>O</i> - <i>trans</i> - <i>p</i> -coumaroyl)- $\beta$ -D-glucopyranoside	<i>E. yuccifolium</i>	[10]
65	kaempferol-3- <i>O</i> -(2,6-di- <i>O</i> - <i>trans</i> - <i>p</i> -coumaroyl)- $\beta$ -D-glucopyranoside	<i>E. yuccifolium</i>	[10]
66	naringenine 7- <i>O</i> - $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- <i>O</i> - $\beta$ -D-glucopyranoside	<i>E. dichotomum</i>	[53]
	<b>Coumarins</b>		
67	umbelliferon	<i>E. biebersteinianum</i>	[89]
68	scopoletin	<i>E. bourgatii</i>	[84]
69	6,7-dimethoxycoumarin	<i>E. creticum</i>	[90]
70	bergaptin	<i>E. biebersteinianum</i>	[89]
71	decursinol	<i>E. biebersteinianum</i>	[89]
72	prantschimgin	<i>E. ilicifolium</i>	[56]
73	deltoin	<i>E. creticum</i>	[56, 91]
74	(+)-marmesin	<i>E. creticum</i> <i>E. ilicifolium</i>	[56, 91]
75	aegelinol benzoate	<i>E. campestre</i>	[55]

Table 2. Contd....

No.	Compound Name	Botanical Source	Reference
76	agasyllin	<i>E. campestre</i>	[55]
77	grandivittin	<i>E. biebersteinianum</i>	[55, 89]
78	aegelinol	<i>E. campestre</i>	[55]
	<b>Phenolics</b>		
79	1- $\beta$ -D-Glucopyranosyloxy-3-methoxy-5-hydroxybenzene	<i>E. creticum</i>	[91]
80	3,4-dihydroxyphenyl caffeate	<i>E. yuccifolium</i>	[10]
81	(4- $\beta$ -D-glucopyranosyloxy)-3-hydroxyphenyl caffeate	<i>E. yuccifolium</i>	[10]
82	R-(+)-rosmarinic acid	<i>E. alpinum</i>	[13]
83	R-(+)-3'-O- $\beta$ -D-glucopyranosyl rosmarinic acid	<i>E. alpinum</i> <i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. maritimum</i> <i>E. campestre</i>	[13]
84	caffeic acid	<i>E. yuccifolium</i>	[10]
85	chlorogenic acid	<i>E. alpinum</i>	[13]
	<b>Steroids</b>		
86	$\beta$ -sitosterol	<i>E. foetidum</i>	[21, 91, 99]
87	$\beta$ -sitosterol 3-O- $\beta$ -D-glucopyranoside	<i>E. creticum</i>	[91]
88	stigmasterol	<i>E. foetidum</i>	[21, 92, 99]
89	stigmasterol 3-O- $\beta$ -D-glucopyranoside	<i>E. dichotomum</i>	[92]
90	campesterol	<i>E. foetidum</i>	[21]
91	brassicasterol	<i>E. foetidum</i> <i>E. agavifolium</i>	[21, Supporting Data I]
92	3 $\alpha$ -cholesterol	<i>E. foetidum</i>	[21]
93	(-)-clerosterol	<i>E. foetidum</i>	[21]
94	$\Delta^5$ -avenosterol	<i>E. foetidum</i>	[21]
95	$\Delta^5$ -avenasterol	<i>E. foetidum</i>	[21]
96	$\Delta^{5,24}$ -stigmastadienol	<i>E. foetidum</i>	[21]
	<b>Acetylenes</b>		
97	falcarinone	<i>E. yuccifolium</i> <i>E. bourgatii</i>	[27, 83, 84]
98	falcarinolone	<i>E. bourgatii</i>	[84]
99	falcarinol	<i>E. yuccifolium</i> <i>E. bourgatii</i>	[27, 84]
100	yuccifolol	<i>E. yuccifolium</i>	[27]
101	1,8-heptadecadiene-4,6-diyne-3,9-diol	<i>E. yuccifolium</i>	[27]
102	(8E)-1,8-Heptadecadiene-4,6-diyne-3,10-diol	<i>E. agavifolium</i>	[Supporting Data I]
103	(Z)-15-hydroxy-9,16-Heptadecadiene-11,13-diyne-8-one	<i>E. agavifolium</i>	[Supporting Data I]
104	(E)-15-hydroxy-9,16-Heptadecadiene-11,13-diyne-8-one	<i>E. agavifolium</i>	[Supporting Data I]
105	(Z)-6-pentyl-2-[2-oxobutyl-(3-ylidene)tetrahydropyran	<i>E. bourgatii</i>	[83, 84]
106	(E)-6-pentyl-2-[2-oxobutyl-(3-ylidene)tetrahydropyran	<i>E. bourgatii</i>	[84]

**Table 3. Other non-Essential oil Compounds Isolated and Identified from *Eryngium***

No.	Name	Botanical Source	Reference
107	2,3,4-trimethylbenzaldehyde	<i>E. varrifolium</i> <i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[46, 82]
108	2,3,6-trimethylbenzaldehyde	<i>E. varrifolium</i> <i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
109	<i>cis</i> -chrysanthenyl acetate	<i>E. planum</i>	[46]
110	<i>cis</i> -chrysanthenyl hexanoate	<i>E. planum</i>	[93]
111	<i>cis</i> -chrysanthenyl octanoate	<i>E. planum</i>	[93]
112	2-angeloyloxymethyl- <i>cis</i> -crotonic acid methyl ester	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
113	2-(2-methyl-butyryloxymethyl)- <i>cis</i> -crotonic acid methylester	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
114	2-isovaleryloxymethyl- <i>cis</i> -crotonic acid methylester	<i>E. bourgatii</i> ( <i>E. amethystinum</i> ) <i>E. serbicum</i>	[82]
115	10-hentriacontanone	<i>E. maritimum</i>	[46]
116	eicosa-8,11-dien-18-ol-2-one	<i>E. creticum</i>	[25]
117	hexadecanoic acid	<i>E. agavifolium</i>	[Supporting Data I]
118	(9Z,11Z)-9,11-octadecadienoic acid, methyl ester	<i>E. agavifolium</i>	[Supporting Data I]
119	(7Z)-7-hexadecen-1-ol, 1-acetate	<i>E. agavifolium</i>	[Supporting Data I]
120	1-kestose	<i>E. planum</i>	[94]
121	sucrose	<i>E. dichotomum</i>	[53, 94]
122	2- <i>O</i> -methyl- $\alpha$ -D-fructofuranose	<i>E. dichotomum</i>	[53]
123	D-glucose	<i>E. planum</i>	[94]
124	D-furanose	<i>E. dichotomum</i>	[53, 94]
125	D-mannitol	<i>E. dichotomum</i> <i>E. creticum</i> <i>E. campestre</i> <i>E. caeruleum</i> <i>E. macrocalyx</i>	[53, 91] [85] [86]
126	D-galactitol	<i>E. creticum</i>	[91]
127	piperidine-2-carboxylic acid	<i>E. maritimum</i>	[88]

23,28-diol (**23**) were isolated and identified from *E. bromeliifolium* [60] and *E. foetidum* [59]. Two glucopyranosyl glycosides of betulinic acid with monosaccharide (**24**) and disaccharide unit (**25**) attached to C-3 were also obtained from *E. bromeliifolium* [61, 62].


## Terpenoids

### Triterpenoids


A total of 8 triterpenoids were isolated and identified from the acid and alkaline hydrolysates of *Eryngium* saponin in a few early investigations (Table 2). All belong to pentacyclic triterpenoids, which are classified as two series of

olean-12-ene and lupine. It is noted that olean-12-ene derivatives are characterized by having multiple hydroxyl groups at C-3, 15, 16, 21, 22, and 28 positions, rarely at C-29 position. These triterpenoids included betulinic acid (**26**) from *E. bromeliifolium* [64], oleanolic acid (**27**) from *E. macrocalyx* (*E. incognitum*) roots [65, 87], R<sub>1</sub>-barringenol (**28**) and A<sub>1</sub>-barringenol (**29**) from the aerial flowering plant parts of *E. maritimum* [63], *E. planum* leaves and roots [95], and the underground parts of *E. caucasicum* (*E. biebersteinianum*) and *E. octophyllum* [66], 22-(2-methyl-2-butenate)-olean-12-ene-3,15,16,22,28-pentol (**30**) from *E. macrocalyx* [65], erynginol A (**32**) from *E. planum* leaves [95], barringtogenol C (**31**) and 22 $\alpha$ -hydroxyerythrodiol (**33**) from *E. maritimum*






- 1 R<sub>1</sub>=H, R<sub>2</sub>=dMA, R<sub>3</sub>=H, R<sub>4</sub>= $\alpha$ -L-rhamnopyranosyl, R<sub>5</sub>=H
- 2 R<sub>1</sub>=OH, R<sub>2</sub>=Ang, R<sub>3</sub>=H, R<sub>4</sub>= $\alpha$ -L-rhamnopyranosyl, R<sub>5</sub>=H
- 3 R<sub>1</sub>=OAc, R<sub>2</sub>=Ang, R<sub>3</sub>=H, R<sub>4</sub>= $\alpha$ -L-rhamnopyranosyl, R<sub>5</sub>=H
- 4 R<sub>1</sub>=OAc, R<sub>2</sub>=dMA, R<sub>3</sub>=H, R<sub>4</sub>= $\alpha$ -L-rhamnopyranosyl, R<sub>5</sub>=H
- 5 R<sub>1</sub>=OH, R<sub>2</sub>=Ang, R<sub>3</sub>=Ac, R<sub>4</sub>= $\alpha$ -L-rhamnopyranosyl, R<sub>5</sub>=H
- 6 R<sub>1</sub>=OH, R<sub>2</sub>=Ang, R<sub>3</sub>=H, R<sub>4</sub>= $\beta$ -D-glucopyranosyl, R<sub>5</sub>= $\alpha$ -L-rhamnopyranosyl
- 7 R<sub>1</sub>=H, R<sub>2</sub>=dMA, R<sub>3</sub>=H, R<sub>4</sub>= $\beta$ -D-glucopyranosyl, R<sub>5</sub>= $\alpha$ -L-rhamnopyranosyl



	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>	R <sub>8</sub>
8	H	OH	H	Glc	H	H	CH <sub>2</sub> OH	Glc
9	H	OH	H	Glc	H	H	CH <sub>2</sub> OH	Gal
10	H	=O	H	Glc	H	H	CH <sub>2</sub> OH	Glc
11	H	OH	H	H	Glc	H	CH <sub>2</sub> OH	Glc
12	OH	OH	OH	Ang	H	Xyl	COOH	H
13	H	OH	OH	Ang	H	Xyl	COOH	H
20	OH	OH	O-Ang	H	H	Ara	COOH	H
14	OH	OH	O-Ang	H	H	Xyl	COOH	H
15	OH	OH	H	Ang	H	Ara	COOH	H
16	OH	OH	H	Ang	H	Xyl	COOH	H
17	OH	OH	O-Ang	Ac	H	Ara	COOH	H
21	OH	OH	O-Ang	Ac	H	Xyl	COOH	H
18	H	OH	O-Ang	Ac	H	Ara	COOH	H
19	H	OH	O-Ang	Ac	H	Xyl	COOH	H




[63]. Barringtonol C (**31**) was also identified being a major component from *E. planum* leaves [95].

### Sesquiterpenes

Four perhydronaphthalene derivatives (Table 2) substituted with one n-propyl were isolated and identified from the hexane/ether extracts of the aerial parts of *E. creticum* grow-

ing in Sinai, Egypt [25] and the etheral extracts of the seeds of *E. giganteum* [46]. The other four sesquiterpenes reported to naturally occur in various essential oils were isolated from the etheral extracts of the seeds of *E. giganteum* [46]. These sesquiterpenes are eryng-9-en-15-al (**34**), 15-hydroxy- $\alpha$ -muurolene (**35**), 15-oxy- $\alpha$ -muurolene (**36**), 15-nor- $\alpha$ -muurolene (**37**), (+)-ledol (**38**), (+)-spathulenol (**39**), germacrene-D (**40**), trans- $\beta$ -farnesene (**41**).



### Monoterpenes

Three monoterpene glycosides of the cyclohexanone type and seven monoterpene aldehyde-esters were identified from seven *Eryngium* species (Table 2). They are include 3-( $\beta$ -D-glucopyranosyloxymethyl)-2,4,4-trimethyl-2,5-cyclohexadien-1-one (42) from *E. campestre* [45] and *E. creticum*, which grows wildly in Jordan [91], 3-( $\beta$ -D-glucopyranosyloxymethyl)-2,4,4-trimethyl-2-cyclohexen-1-one (43) from *E. campestre* [45], 5-[( $\beta$ -D-glucopy-

ranosyloxy)methyl]-4-hydroxy-4-[(1*E*,3*S*)-3-hydroxy-1-butenyl]-3,5-dimethyl-2-cyclohexen-1-one (44) from the aerial parts of *E. dichotomum* from Tunisian [53], isoferulyl senecioate (45) from leaves of *E. variifolium* [46], (-)-2,4,4-trimethyl-3-formyl-2,5-cyclohexadienyl angelate (46) from seeds of *E. paniculatum* [47], *O*-[2-angeloyloxymethyl-*cis*-crotonoyl]-ferulol (47), *O*-[2-angeloyloxymethyl-*cis*-crotonoyl]-isoferulol (48), *O*-[2-(2-methyl-butyryloxymethyl)-*cis*-crotonoyl]-ferulol (49), *O*-[2-(2-methyl-butyryloxymethyl)-*cis*-crotonoyl]-isoferulol (50), and *O*-[2-isovaleryloxy-

methyl-*cis*-crotonoyl]-isoferulol (**51**) from the roots of *E. bourgatii* (*E. amethystinum*) and *E. serbicum* [82].

## Phenolic Compounds

### Flavonoids

Flavonoids are also one of the main components of the genus *Eryngium*. A total of 15 flavonoids including flavonols (**52-65**), and flavanone (**66**) were isolated and identified from *Eryngium* species (Table 2). The glycosides of quercetin (**52**) [91] and kaempferol with an oligosaccharide chain at C-3 or C-7, or two oligosaccharide chains at C-3 and C-7 are types of flavonol glycosides.

Chemical investigation on the aerial parts of *E. campestre* resulted in the isolation and structural determination of 8 flavonol glycosides [50, 52]. These compounds are quercitrin (**53**), isoquercitrin (**54**), rutin (**55**), astragalin (**57**), kaempferol 7-*O*- $\alpha$ -L-rhamnopyranoside (**58**), kaempferol 3,7-di-*O*- $\alpha$ -L-rhamnopyranoside (**60**), Kaempferol 3-*O*- $\beta$ -D-(2'-*p*-*E*-hydroxycinnamoyl)-glucopyranoside (**62**), and kaempferol 3-*O*- $\beta$ -D-(2'-*p*-*Z*-hydroxycinnamoyl)-glucopyranoside (**63**). Kaempferol 3- $\beta$ -D-glucopyranosyl-7-*O*- $\alpha$ -L-rhamnopyranoside (**59**), astragalin, and isoquercitrin was identified as the major flavonoid constituents from the aerial parts of *E. maritimum* [49]. Three investigations led to the isolation and identification of two flavonoid glycosides Kaempferol-3-*O*-(6-*O*- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranoside (**61**) and **60** from the leaves of *E. planum* [96-98]. A new compound, kaempferol-3-*O*-(2-*O*-*trans*-*p*-methoxycoumaroyl)-6-*O*-*trans*-*p*-coumaroyl)- $\beta$ -D-glucopyranoside (**64**), and a known flavonoid glycoside kaempferol-3-*O*-(2,6-di-*O*-*trans*-*p*-coumaroyl)- $\beta$ -D-glucopyranoside (**65**) were recently isolated and identified from the whole plants of *E. yuccifolium* [10]. Additionally, Compound **55** was also found from the aerial parts of *E. octophyllum* [51], compound **59** from the aerial parts of *E. macrocalyx* [51], and compounds **54** and **60** from the leaves of *E. giganteum* [54].

Luteolin 7-*O*- $\beta$ -D-glucopyranoside (**56**) and naringenin 7-*O*- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)-*O*- $\beta$ -D-glucopyranoside (**66**) were also identified from *E. campestre* [52] and the aerial parts of *E. dichotomum* [53].

### Coumarins

Simple coumarins and linear furanocoumarins were also found from *Eryngium* species (Table 2). These coumarins are scopoletin (**68**) from the roots of *E. bourgatii* [84]; 6,7-dimethoxycoumarin (**69**), deltoin (**73**), and (+)-marmesin (**74**) from *E. creticum* [90, 91]; umbelliferon (**67**), bergaptin (**70**), decursinol (**71**), grandivittin (**77**) from *E. biebersteini-anum* [89]; prantschimgin (**72**), deltoin, and (+)-marmesin from Et<sub>2</sub>O extracts of *E. ilicifolium* [56]; and aegelinol benzoate (**75**), agasyllin (**76**), aegelinol (**78**), and grandivittin from the roots of *E. campestre* [55].

### Other Phenolic Compounds

1- $\beta$ -D-glucopyranosyloxy-3-methoxy-5-hydroxybenzene (**79**) was yielded from *E. creticum*, which grows wildy in Jordan [91]. Three phenolic compounds, including two new phenyl caffeates, 3,4-dihydroxyphenyl caffeate (**80**) and (4- $\beta$ -D-glucopyranosyloxy)-3-hydroxyphenyl caffeate (**81**), together with known compound caffeic acid (**84**) were isolated and identified from *E. yuccifolium* [10].

In addition, the isolation of antioxidative substances from the root extracts of *E. alpinum* allowed the identification of R-(+)-3'-*O*- $\beta$ -D-glucopyranosyl rosmarinic acid (**83**), a new rosmarinic acid derivative, together with two known compounds R-(+)-rosmarinic acid (**82**), and chlorogenic acid (**85**) [13]. Moreover, R-(+)-3'-*O*- $\beta$ -D-glucopyranosyl rosmarinic acid (**83**) was found that is unstable in a MeOH/H<sub>2</sub>O (1/1, v/v) solution, of which one-third quantity was degraded after 11 h at room temperature. In the chemotaxonomic study analyzed by HPLC, R-(+)-rosmarinic acid and R-(+)-3'-*O*- $\beta$ -D-glucopyranosyl rosmarinic acid were detected in all analyzed 13 *Eryngium* species, except *E. giganteum*, which was devoid of R-(+)-3'-*O*- $\beta$ -D-glucopyranosyl rosmarinic acid, and distinct concentration variations of **83** were observed in *E. bourgatii* (*E. amethystinum*), *E. maritimum*, and *E. campestre* [13].

### Steroids


Eleven steroids have been identified from *Eryngium* species (Table 2). These compounds include  $\beta$ -sitosterol (**86**), stigmasterol (**88**), campesterol (**90**), brassicasterol (**91**), 3 $\alpha$ -cholesterol (**92**), (-)-clerosterol (**93**),  $\Delta^5$ -avenosterol (**94**),  $\Delta^7$ -avenasterol (**95**), and  $\Delta^{5,24}$ -stigmastadienol (**96**) from the hexane extracts of the leaves of *E. foetidum* [21, 99]. Brassicasterol (**91**) was also isolated and identified from the EtOH extract of the whole plant of *E. agavifolium* (Supporting Data I). A comparison study indicated that all of steroids in the leaves presented also in the stems with different quantities [99]. Stigmasterol,  $\beta$ -sitosterol, and their glycosides (**87** and **89**) were isolated and identified from the aerial parts of *E. dichotomum* grown in Tunisia [92] and *E. creticum* growing wildy in Jordan [91], respectively.

### Acetylenes

*Eryngium* contains also polyacetylenes (Table 2). Yuccifolol (nonadeca 1,11-diene 4,6,8-triyn 3,10-diol, **100**), a novel polyacetylene, was isolated and identified from the hexane/ether extracts of the aerial parts of *E. yuccifolium*, together with the known polyacetylenes, falcarinone (**97**), falcarinol (**99**) and heptadeca 1,8-diene 4,6-diyne 3,9-diol (**101**) [27]. (8*E*)-1,8-heptadecadiene-4,6-diyne-3,10-diol (**102**), (Z)-15-hydroxy-9,16-heptadecadiene-11,13-diyne-8-one (**103**), and (E)-15-hydroxy-9,16-heptadecadiene-11,13-diyne-8-one (**104**) were isolated from the EtOH extract of the whole plant of *E. agavifolium* (Supporting Data I). The roots of *E. bourgatii* afforded, in addition, to falcarinone, falcarinolone (**98**), and falcarinol, the Z and E isomers of 6-pentyl-2-[2-oxobutin-(3)-yliden]tetrahydropyran (**105** and **106**). The new Z isomer of 6-pentyl-2-[2-oxobutin-(3)-yliden]tetrahydropyran is unstable as it easily converts into the E isomer [83, 84].

### Miscellaneous


There are 18 other classes of compounds obtained from some species of *Eryngium* L. (Table 3) These compounds include two trimethylbenzaldehydes (**107** and **108**) from the leaves of *E. varrifolium* [46], *E. bourgatii* (*E. amethystinum*) and *E. serbicum* [82], three *cis*-chrysanthenyl esters (**109-111**) from the flowers and seeds of *E. planum* [46, 93], three 2-substituents derivatives of *cis*-crotonic acid methyl ester (**112-114**) from *E. bourgatii* (*E. amethystinum*) and *E. ser-*



*bicum* [82], two long-chain aliphatic ketones (**115** and **116**) from the leaves of *E. maritimum* [46] and *E. creticum* [25], and three long-chain aliphatic acids (**117-119**) from the EtOH extract of the whole plant of *E. agavifolium* (Supporting Data I). From a chemotaxonomic point of view, 10-hentriacontanone (**115**) appears to be a good chemical marker of *E. maritimum*. Analysis of several *E. maritimum* collections of different origins always showed the presence of this ketone. However, this metabolite has also been found

in two other *Eryngium* species, namely *E. bourgatii* and *E. campestre* [46].

In addition, sucrose (**121**), furanose (**124**) and its 2-methyl analog (**122**) were found from the aerial parts of *E. dichotomum* in Tunisian [53]; D-glucose (**123**) and a nonreducible trisaccharide, 1-kestose (**120**), from *E. planum* roots [94]; two polyols, D-mannitol (**125**) and D-galactitol (**126**), from *E. dichotomum* [53], *E. creticum* [91], *E. campestre*



[85], *E. caeruleum* and *E. macrocalyx* [86]; and a piperidine-2-carboxylic acid (**127**) from *E. maritimum* [88].


## BIOLOGICAL AND PHARMACOLOGICAL ACTIVITIES

The *in vitro* and *in vivo* biological activities of *Eryngium* extracts or isolates (including both essential oils and non-essential oil compounds) are summarized in Tables 4 and 5.

### Cytotoxicities

Eryngiosides A-C, E, F, H-J, L, 21β-angeloyloxy-3β-[β-D-glucopyranosyl-(1→2)]-[β-D-xylopyranosyl-(1→3)]-β-D-glucuronopyranosyloxyolean-12-ene-15α, 16α, 22α, 28-tetrol (**20**), and Saniculasaponin III isolated from the whole plants of *E. yuccifolium* Michx [10] were evaluated for their

ability to inhibit human DNA topoisomerase I (TOP1) and II (TOP2) activity and cytotoxicity against PANC-1, A549, PC-3, HL-60, and MRC-5 cell lines (Table 6). None of eryngiosides demonstrated any TOP1 and TOP2 inhibitory activities. Three eryngiosides (eryngioside J, eryngioside L, and saniculasaponin III) showed moderate cytotoxicity against all cell lines after 48 h of incubation. The GI<sub>50</sub> values vary from 3.54 μM to 16.02 μM. These eryngiosides were most effective against human lung cancer cell lines A-549 and normal cell lines MRC-5, and markedly inhibited the growth of pancreas cancer cell lines PANC-1. *Eryngium* saponin **20** also exhibited moderate cytotoxicity against A549, PC-3, HL-60, and MRC-5 cell lines with GI<sub>50</sub> values of 7.34-9.38 μM while no toxicity against the human pancreas cancer cells PANC-1 [81]. Especial, Eryngioside H and I exhibited potent and highly selective inhibition against four



human tumor cells but almost no cytotoxicity against normal human cells (Figs. 1 and 2, and see Supporting Data II). Compounds 1-4, and 7 isolated from the roots of *E. campstre* showed a weak cytotoxic activity, with IC<sub>50</sub> between 40 and 100 μg/mL, against HCT 116 and HT-29 human tumor cell lines by MTT assay [15].

Ethanol extracts from fruits of *E. planum* were shown to display highly significant in inducing apoptosis in two human leukemic cell lines C8166 (96%) and J45 (89%) after 24 h incubation in concentration of 300 μg/mL. Lower level of apoptotic cells was observed in HL-60 (49%) and ML-1 (42%). The tested human leukaemia cell lines included human acute myeloblastic leukaemia (ML-1), human acute T

cell leukaemia (J-45.01), human eosinophilic leukaemia (EOL), human caucasian promyelocytic leukaemia (HL-60), human T cell leukaemia lymphoblast (1301), human T cell leukaemia (C-8166), human myeloma (U-266B1), human Caucasian normal B cell (WICL), and human T cell (H-9) [101].

#### Anti-Mutagenic Activity

An anti-mutagenic potential of *E. creticum* was evidenced on MNNG induced mutagenicity using rat hepatocytes. The study was focused on ethanolic extracts, while the extracts had no effect on cytotoxicity indicators such as necrosis and apoptosis. The effects obtained can be attributed

**Table 4. Biological Activities of *Eryngium* as Revealed by *In Vitro* Studies**

Bioactivities	Description	Bioactive Agents	Reference
Cytotoxicities	Exhibited moderate or weak cytotoxicities against human pancreas, prostate, lung, leukemic, colon cancer cell lines and normal human lung fibroblast cell line	Saponin Compounds	[15, 81, Supporting Data II]
	Ethanol extracts induced apoptosis in human leukemic cell lines	Ethanol Extracts	[101]
Anti-mutagenic Activity	Exhibited an anti-mutagenic potential on MNNG mutagenicity using rat hepatocytes	Ethanol Extracts	[102]
Anti-inflammatory Activities	Extracts Inhibited cytokine-stimulated, iNOS-dependent synthesis of nitric oxide in murine endothelial cells, without affecting cell viability	Extracts	[103]
	Reduced myeloperoxidase activity strongly in the inflamed tissue of the acute model	Polar Sterols	[21]
Anti-snake and Scorpion Venoms Effects	Aqueous extracts of the leaves and roots inhibited the hemolytic activity of the snake and scorpion venoms	Aqueous Extracts	[24]
	Ethanol extracts of the leaves and roots enhanced RBC hemolysis on red blood cells	Ethanol Extracts	[24]
	Inhibited the contraction of isolated tracheal smooth muscle of rabbits and	Aqueous Extract	[106]
Antibacterial, Antifungal, and Antimalarial Activities	Exhibited selective antibacterial activity against <i>Salmonella</i> species and the <i>Erwinia</i> genus of bacteria	Eryngial	[108, 109]
	Exhibited activity against parasitic trypanosomes, nematodes, fungi and bacteria in humans and other mammals	Eryngial	[109]
	Exhibited antifungal activity with MIC values of 0.16-0.32 µg/mL against several dermatophyte species	Volatile Extracts	[110]
	Showed antimycotic activity	Saponin Mixture Extracts	[111, 113, 114]
	Showed antimicrobial activities against 12 bacterial and yeast strains	Chloroformic Fractions	[112]
	Showed antiplasmodial activity against <i>Plasmodium falciparum</i> with IC <sub>50</sub> of 25 µg/mL and <i>P. gallinaceum</i> which infects chickens	Extracts	[109]
Antioxidant Activities	Exhibited antioxidant activity in <i>in vitro</i> assays	Rosmarinic acid analogs Extracts	[13, 115-121]
	Exhibited different levels of antioxidant activity six <i>in vitro</i> assays at both flowering and non-flowering stage	Methanol Extracts	[26, 122]
	Exhibited weak radical scavenging activity, low total phenol content	Extracts	[112]
	Inhibited iron-fructose-phosphate-induced lipid peroxidation in lecithin liposome and linoleic acid emulsion systems	Ethanol Extracts	[123]
Other Activities	Exhibited the inhibition in the velvetleaf germination	(-)-2,4,4-trimethyl-3-formyl-2,5-cyclohexadienyl angelate	[47]
	Exhibited anthelmintic activity against <i>Strongyloides stercoralis</i>	Eryngial	[109]

to a direct antimutagenic activity and an increased recovery at the chromosomal level [102].

### Anti-Inflammatory Activities

Extracts obtained from the root and aerial parts of various *Eryngium* species are used as folk remedy worldwide for the treatment of various inflammatory disorders. Recently, in a antiinflammatory activity screening on extracts of 121 plants typical for the traditional Mediterranean diet, *E. campestre* L. decreased nitric oxide and TNF-alpha synthesis in the murine endothelial cells of monocyte origin activated with LPS, decreased cytokine or LPS-stimulated iNOS mRNA levels in both cell types [103]. It has been reported that ethanol extracts either from the aerial parts or roots of 7 *Eryn-*

*gium* species growing in Turkey showed remarkably anti-inflammatory and antinociceptive activity in mice. Especially, the aerial parts and roots of *E. maritimum* and *E. kotschyi* were found to possess most promising activities without including any apparent gastric damage [22]. More detailed research showed the antiinflammatory effect of *E. maritimum* may be produced by a sub-alkaline fraction [23]. Saponin mixtures isolated from *E. planum* prevented inflammatory responses when injected into rats at concentration of 0.5 mg/kg, but oral administration at same dose had no antiinflammatory effect [104].

The hexane extracts from the leaves of *E. foetidum* can reduce the edema, induced by 12-0-tetradecanoylphorbol acetate (TPA) in the mouse, in a similar proportion in acute

**Table 5. Biological Activities of *Eryngium* as Revealed by *In Vivo* Studies**

Bioactivities	Animal Model	Description	Bioactive Agents	Reference
Anti-inflammatory Activities	Mice	Inhibited carrageenan-induced hind paw oedema and TPA-induced ear oedema tests	Ethanol Extracts	[22]
	Mice	Showed antinociceptive activity in the p-benzoquinone-induced writhing test	Ethanol Extracts	[22]
	Mice	Reduced the auricular oedema in acute and chronic assay induced by 12-0-tetradecanoylphorbol acetate (TPA)	Polar Sterols	[21]
	Rats	Prevented inflammatory responses to s.c. injections of nucleic acid Na salt or ovalbumin into the hind-paw	Saponin Mixtures	[104]
	Rats	Inhibited the carrageenan-induced oedema in paw given orally	Decoctions	[105]
	Rats	Induced the number of abdominal writhing provoked by acetic acid as the pain stimulus	Decoctions	[105]
Anti-Scorpion Venoms Effects	Guinea pigs	prolonged the life from 20 min to 8 hr injected by Jordanian <i>Leiurus quinquestriatus</i> scorpion venom	Aqueous Extracts	[106, 107]
Antihyperglycemic Effects	Rats	Reduced blood glucose concentration given orally in normoglycemic and streptozotocin -induced models	Decoctions	[17]
	Rats	Exhibited substantial acute antihyperglycemic activities despite lacking any favorable <i>in vitro</i> effectiveness	Aqueous Extracts	[124]
	Rats	Showed no effect in the level of glucose of the normoglycemic and streptozotocin -induced diabetic, and normal models when given a single oral dose	Extracts	[109]
Other Activities	Rats	Exhibited anti-convulsant activity at a concentration of 110 g/250 mL induced by picrotoxin	Extracts	[109, 127]
	Rats	Enhanced the permeation of piroxicam across rat skin	Essential Oil	[125, 126]

and chronic assay. Myeloperoxidase activity was also strongly reduced in the acute, but not the chronic model. Although stigmasterol (**88**) yielded from the hexane extracts exerted a significant topical antiinflammatory activity, by itself it could not account for the overall effects observed for the total phytosterols [21]. Further study indicated that a polar sterol should be responsible for the so-called medicinal property of the plant [99]. Additionally, a decoction from the leaves of *E. foetidum* exhibited significantly dose-dependent anti-inflammatory activity by inhibiting the carrageenan-induced oedema in rat paw. However, oral administration was less active than topical administration. This decoction also potentially decreased the number of abdominal writhings provoked by acetic acid as the pain stimulus [105].

#### Anti- Snake and Scorpion Venoms Effects


Traditionally, many *Eryngium* species were used to prevent and treat snake bites and scorpion stings [10]. Modern pharmacologic investigations are only limited to *E. creticum*. *E. creticum* has been used in folk medicine in Jordan as a

remedy for scorpion stings in the rural areas and as a hypoglycemic agent [17]. Aqueous extracts of both fresh and dried roots as well as fresh leaves of *E. creticum* gave 100% or a higher percentage inhibition of the hemolytic activity of the snake and scorpion venoms. However, ethanol extract of the leaves and roots enhanced red blood cell (RBC) hemolysis rather than inhibiting venom activities on RBC [24]. Aqueous extract of the roots inhibited the contraction of isolated tracheal smooth muscle of rabbits and guinea pigs caused by *Leiurus quinquestriatus* scorpion venom, and also prolonged the life of guinea pigs from 20 min to 8 h when injected by a Jordanian *L. quinquestriatus* scorpion venom [106, 107].

#### Antibacterial, Antifungal, and Antimalarial Activities

The aerial plant parts of *E. foetidum* exhibited selective antibacterial activity against *Salmonella* species and the *Erwinia* genus of bacteria [108, 109]. A fraction of the essential oil rich in eryngial is the subject of a US patent application for its effectiveness against parasitic trypanosomes, nematodes, fungi and bacteria in humans and other mammals [109]. The volatile extracts of *E. duriaei* subsp. *juresianum*





**Table 6. Inhibitory Activity Against DNA Topoisomerases and Cytotoxicity Against Human Tumors of Some Oleanane-type Triterpenoids and Triterpenoid Glycosides**

No.	Name	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>	Activity (IC <sub>50</sub> ) (μM)		Cytotoxicity (GI <sub>50</sub> ) (μM)				
									TOP1	TOP2	A549	PC-3	HL-60	PANC-1	MRC-5
1	Eryngioside J	OH	OH	O-Ang	Ac	Ara	COOH	H	(-)	(-)	4.3±0.34	5.08±0.38	7.36±0.28	11.45±0.75	5.61±1.27
2	Saniculasaponin III	OH	OH	O-Ang	Ac	Xyl	COOH	H	(-)	(-)	4.73±0.21	8.51±0.71	7.11±0.55	16.02±0.61	3.54±0.34
3	Eryngioside L	H	OH	O-Ang	Ac	Xyl	COOH	H	(-)	(-)	8.35±2.89	11.32±1.60	8.41±1.84	13.0±1.27	8.71±1.45
4	21β-angeloyloxy-3β-[β-D-glucopyranosyl-(1→2)]-[β-D-xylopyranosyl-(1→3)]-β-D-glucuronopyranosyloxy-olean-12-ene-15α, 16α, 22α, 28-tetrol	OH	OH	O-Ang	H	Ara	COOH	H	(-)	(-)	7.34±0.46	7.99±0.29	9.38±1.48	(-)	8.51±1.27
5	Eryngioside F	H	OH	OH	Ang	Xyl	COOH	H	(-)	(-)	(-)	(-)	(-)	(-)	(-)
6	Eryngioside E	OH	OH	OH	Ang	Xyl	COOH	H	(-)	(-)	(-)	(-)	(-)	(-)	(-)
7	Eryngioside H	OH	OH	H	Ang	Ara	COOH	H	(-)	(-)	1.0±0.15	2.12±0.11	3.65±1.00	12.71±2.19	5.61±0.71
8	Eryngioside I	OH	OH	H	Ang	Xyl	COOH	H	(-)	(-)	1.2±0.16	2.71±0.06	2.19±0.68	7.06±0.94	4.59±0.36
9	Eryngioside A	H	OH	H	Glc	H	CH <sub>2</sub> OH	Glc	(-)	(-)	(-)	(-)	(-)	(-)	(-)
10	Eryngioside B	H	OH	H	Glc	H	CH <sub>2</sub> OH	Gal	(-)	(-)	(-)	(-)	(-)	(-)	(-)
11	Eryngioside C	H	O	H	Glc	H	CH <sub>2</sub> OH	Glc	(-)	(-)	(-)	(-)	(-)	(-)	(-)

Notes: *DNA Topoisomerase Inhibitory Activity*: For active compounds, IC<sub>50</sub> (mean ± S.D.) refers to the concentration required to inhibit 50% of TOP1 activity. (-) indicates that the compound is inactive (negative at 312 μM or with IC<sub>50</sub> >250 μM). *Cytotoxicity*: GI<sub>50</sub> (mean ± S.D.) refers to the concentration required to have 50% cell-growth inhibition; (-) indicates that the compound is inactive at 25 μM

showed antifungal activity with MIC values of 0.16-0.32 μL/mL against several dermatophyte species (*Trichophyton mentagrophytes*, *T. rubrum*, *Epidermophyton floccosum*; *T. verrucosum*, *T. mentagrophytes var interdigitale*, *Microsporum canis* and *M. gypseum*) [110]. In an *in vitro* antimycotic activity screening against 8 phytopathogenic fungi, *E. creticum* showed more than 95% inhibition of spore germination in at least two fungi [111]. Two extracts from *E. maritimum* L. showed antimicrobial activities against 12 bacterial and yeast strains. The result indicated that chloroformic fractions were generally more active than methanolic ones [112]. A saponin mixture from dried leaves of *E. planum* also showed antimycotic effect [113, 114].

Despite the claims of traditional antimalarial use, *E. foetidum* showed lowly *in vitro* antiparasmodial activity against *Plasmodium falciparum* with IC<sub>50</sub> of 25 μg/mL. Interestingly, in the screening of the aqueous extract of the entire plant against various species of *Plasmodium*, activity was only reported against *P. gallinaceum* which infects chickens thereby suggesting another possible veterinary use [109].


### Antioxidant Activities

There is currently an upsurge of interest in phytochemicals as new sources of natural antioxidants. In several *in vi-*


*tro* antioxidant activity screening, many *Eryngium* species used as edible plants [115] and/or herbs in Thailand [116], Jordanian origin [117], Vietnamese [118], European [119], Sardinia [120], and Iran [121] have been demonstrated to have the antioxidant activity in tested models.

The roots of *E. alpinum* were shown to have highly antioxidant activity toward the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical in a TLC autographic assay. By further bioassay fractionation, compounds **82**, **83** and **85** were found to respond for the activity. Related to the activity of ascorbic acid, the antioxidant activity of **82** was almost one-fold higher than **83**, and similar to **84**. The results are consistent with published reports indicating that the position and/or the number of glycosyl groups present in the molecule plays a significant part in the antioxidant activity [13].

Methanol extracts of leaves and inflorescence of *E. caucasicum* at flowering stage, which was found recently as a new cultivated vegetable plant in home gardens in northern Iran, were investigated for their antioxidant activities employing six *in vitro* assay systems. Extracts exhibited different levels of antioxidant activity in all the models studied. Extracts showed very good scavenging activity of H<sub>2</sub>O<sub>2</sub> with IC<sub>50</sub> of 25.5 mg/mL for leaves and 177.2 mg/mL for inflorescence, respectively; IC<sub>50</sub> for DPPH radical-scavenging



**Fig. (1).** Effects of Eryngioside H and I on the proliferation of human non-small cell lung cells (A549).



**Fig. (2).** Effects of Eryngioside H and I on the proliferation of human pancreatic cancer cells (PANC-1).

activity was 0.15 for leaves and 0.39 mg/mL for inflorescence; leaves extract exhibited better  $\text{Fe}^{2+}$  chelating ability ( $\text{IC}_{50} = 0.25$  mg/mL) than that of EDTA ( $\text{IC}_{50} = 18$   $\mu\text{g}/\text{mL}$ ) [26]. The research group also reported a good antioxidant activity of leaves of *E. caucasicum* at non-flowering stage using the same assay systems. However, these are so different from flowering stages. The leaf extracts at non-flowering stage showed stronger NO scavenging and peroxidation inhibition, and very less scavenging activity of  $\text{H}_2\text{O}_2$ , reducing powers and  $\text{Fe}^{2+}$  chelating ability than leaves and inflorescence at flowering stage [122].

*Eryngium maritimum* exhibited weak radical scavenging activity ( $\text{IC}_{50} = 0.28$  mg/mL), low total phenol content (16.4 mg/g), as well as relatively strong total antioxidant activities (from 32.7 to 48.6 mg/g) [112]. Ethanol extract of *E. billardieri* significantly inhibited iron-fructose-phosphate-induced lipid peroxidation in lecithin liposome and linoleic acid emulsion systems [123].

### Antihyperglycemic Effects

A research indicated that an aqueous decoction from *E. creticum* caused significant reductions in blood glucose concentration when given orally in normoglycemic and streptozocin-hyperglycemic rats [17]. The aqueous extracts of *E. creticum* exhibited substantial acute antihyperglycemic ac-

tivities in starch-treated rats, despite lacking any favorable *in vitro* effectiveness [124]. In a preliminary oral glucose-tolerance evaluation, a single (acute) oral dose of the leaf extract from *E. foetidum* did not cause significant reduction in the level of glucose of the models tested [109].

In addition, some other bioactivity and pharmacological properties of *Eryngium* also report. (-)-2,4,4-trimethyl-3-formyl-2,5-cyclohexadienyl angelate (**46**), only active compound isolated from the hexane extract of seeds from *E. paniculatum* was found to inhibit germination of velvetleaf [47]. The essential oil of *E. bungei* and *E. caeruleum* at the 5.0% concentration provided an almost 9.17-fold and 8.56-fold increase in permeability coefficients of piroxicam across rat skin [125, 126]. Pharmacological studies of the aerial plant parts of *E. foetidum* L. included also anthelmintic activity due to eryngial and anti-convulsant activity at a concentration of 110 g/250 mL induced by picrotoxin in the respective models [109, 127]. *Eryngium foetidum* is one of four plants used in a Japanese patent for having developed a skin-whitening agent [109].

### CONCLUSIONS

*Eryngium* has been cultivated as ornamental, vegetable, and medicinal crops. However, phytochemistry and pharmacological properties of most of the 250 species remain unex-

plored. From the 23 species, at least 127 compounds have been isolated and identified. These are primarily non-essential oil compounds such as terpenoids, triterpenoid saponins, flavonoids, coumarins, polyacetylenes, and steroids. These 23 species represent nine to 11 sections of *Eryngium* in both Old and New Worlds. Initial data indicated that triterpenoid saponins, terpenoids, coumarins may be restricted to certain sections. Because the chemical investigations for the most of these species except *E. campestre* and *E. yuccifolium* are preliminary and incomplete, chemotaxonomical significance of these compounds will not be revealed until extensive investigations including more species.

*Eryngium* extracts or isolates have shown *in vitro* and *in vivo* activities such as cytotoxicity against various human tumor cell lines, anti-inflammatory, anti-snake and scorpion venoms, antibacterial, antifungal, and antimalarial, antioxidant, and antihyperglycemic effects. There are no clinical data related to the numerous ethno medicinal uses. The molecular mechanisms of bioactivities (particularly cytotoxicity and anti-snake and scorpion venoms) of *Eryngium* isolates remain elusive. Also, chemical isolation and modification and animal tests of polyhydroxylated triterpenoid saponins may provide interesting lead for cancer drug development.

#### ABBREVIATIONS

A549	= Human lung adenocarcinoma epithelial cell line
Ang	= Angeloyl
Ac	= Acetyl
dMA	= $\beta,\beta$ -dimethylacryloyl
DPPH	= 2,2-diphenyl-1-picrylhydrazyl
EDTA	= Ethylenediaminetetraacetic acid
HL-60	= Human promyelocytic leukemia cell line
HCT 116	= Human colon adenocarcinoma epithelial cell line
HT-29	= Human colon adenocarcinoma epithelial cell line
iNOS	= Inducible nitric oxide synthase
LPS	= Lipopolysaccharide
MNNG	= N-methyl-N'-nitro-N-nitroso-guanidine
MRC-5	= Human normal lung cell line
mRNA	= Messenger ribonucleic acid
PANC-1	= Human pancreatic carcinoma, epithelial-like cell line
PC-3	= Human prostate adenocarcinoma epithelial cell line
RBC	= Red blood cell
SAR	= Structure-activity relationship
TOP1	= DNA topoisomerase I
TOP2	= DNA topoisomerase II
TNF	= Tumor necrosis factor
TPA	= 12-O-tetradecanoylphorbol acetate

#### CONFLICT OF INTEREST

None Declared.

#### ACKNOWLEDGEMENTS

This work was funded by the USDA grant 2010-38928-21678.

#### SUPPLEMENTARY MATERIAL

Supplementary material is available on the publishers Web site along with the published article.

#### REFERENCES

- [1] Wörz, A. On the distribution and relationships of the South-West Asian species of *Eryngium* L. (Apiaceae-Saniculoideae). *Turk. J. Bot.*, **2004**, 28, 85-92.
- [2] Wörz, A. A new subgeneric classification of the genus *Eryngium* L. (Apiaceae, Saniculoideae). *Bot. Jahrb. Syst.*, **2005**, 126(2), 253-259.
- [3] Calviño, C. I.; Martínez, S. G.; Downie, S. R. The evolutionary history of *Eryngium* (Apiaceae, Saniculoideae): rapid radiations, long distance dispersals, and hybridizations. *Mol. Phylogenet. Evol.*, **2007**, 46 (3), 1129-1150.
- [4] Rodríguez-Gacio, M. C.; De Justús, J.; María, R.; María, H. Genetic diversity among genotypes of *Eryngium viviparum* (Apiaceae): a plant threatened throughout its natural range. *Bot. J. Linnean Soc.*, **2009**, 159 (2), 237-244.
- [5] Gaudeul, M.; Taberlet, P.; Till-Bottraud, I. Genetic diversity in an endangered alpine plant, *Eryngium alpinum* L. (Apiaceae), inferred from amplified fragment length polymorphism markers. *Mol. Ecol.*, **2001**, 9 (10), 1625-1637.
- [6] Dolan, R. W.; Yahr, R.; Menges, E. S.; Halfhill, M. D. Conservation implications of genetic variation in three rare species endemic to Florida rosemary scrub. *Am. J. Bot.*, **1999**, 86, 1555-1562.
- [7] Boice, L. P. Managing endangered species on military lands. *Endang. Sp. UPDATE*, **1996**, 13 (7 & 8), 1-6.
- [8] Wolff, H. In: *Das Pflanzenreich*, Engler, A., Ed. Wilhelm Engelmann: Leipzig and Berlin, **1913**, vol. 228, pp. 1-305.
- [9] Wörz, A.; Diekmann, H. Classification and evolution of the genus *Eryngium* L. (Apiaceae-Saniculoideae): results of fruit anatomical and petal morphological studies. *Plant Divers. Evol.*, **2010**, 128 (3-4), 387-408.
- [10] Zhang, Z. Z.; Li, S. Y.; Ownby, S.; Wang, P.; Yuan, W.; Zhang, W. L.; Beasley, R. S. Phenolic compounds and rare polyhydroxylated triterpenoid saponins from *Eryngium yuccifolium*. *Phytochemistry*, **2008**, 69, 2070-2080.
- [11] Ekpong, B.; Sukprakarn, S. Harvest stages and umbel order contribution on erylgo (*Eryngium foetidum* L.) seed yield and quality. *Kasetsart J.*, **2006**, 40, 273-279.
- [12] Khoshbakht, K.; Hammer, K.; Pistrick, K. *Eryngium caucasicum* Trautv. cultivated as a vegetable in the Elburz Mountains (Northern Iran). *Genet Resour. Crop Evol.*, **2006**, 54 (2), 445-448.
- [13] Le Claire, E.; Schwaiger, S.; Banaigs, B.; Stuppner, H.; Gafner, F. Distribution of a new rosmarinic acid derivative in *Eryngium alpinum* L. and other Apiaceae. *J. Agric. Food Chem.*, **2005**, 53, 4367-4372.
- [14] Kartal, M.; Mitaine-Offer, A. C.; Abu-Asaker, M.; Miyamoto, T.; Calis, I.; Wagner, H.; Lacaille-dubois, M. A. Two new triterpene saponins from *Eryngium campestre*. *Chem. Pharm. Bull.*, **2005**, 53 (10), 1318-1320.
- [15] Kartal, M.; Mitaine-Offer, A. C.; Paululat, T.; Abu-Asaker, M.; Wagner, H.; Mirjolet, J. F.; Guilbaud, N.; Lacaille-Dubois, M. A. Triterpene saponins from *Eryngium campestre*. *J. Nat. Prod.*, **2006**, 69, 1105-1108.
- [16] Baytop, T. *Therapy with Medicinal Plants in Turkey-Past and Present*. 2nd ed.; Nobel Tip Basimevi: Istanbul **1999**.
- [17] Jaghabir, M. Hypoglycemic effects of *Eryngium creticum*. *Arch. Pharm. Res.*, **1991**, 14 (4), 295-297.
- [18] Goleniowski, M. E.; Bongiovanni, G. A.; Palacio, L.; Nunez, C. O.; Cantero, J. J. Medicinal plants from the "Sierra de Comechingones", Argentina. *J. Ethnopharmacol.*, **2006**, 107, 324-341.

- [19] Bencao, E. C. O. Z. *Zhonghua Bencao*. Shanghai Science and Technology Press: Shanghai, **1999**, Vol. 5.
- [20] Moerman, D. E. *Native American Medicinal Plants: An Ethnobotanical Dictionary*. Timber Press: Portland, **2009**.
- [21] García, M. D.; Sáenz, M. T.; Gómez, M. A.; Fernández, M. A. Topical antiinflammatory activity of phyosterols isolated from *Eryngium foetidum* on chronic and acute inflammation models. *Phytother. Res.*, **1999**, *13* (1), 78-80.
- [22] Kupeli, E.; Kartal, M.; Aslan, S.; Yesilada, E. Comparative evaluation of the anti-inflammatory and antinociceptive activity of Turkish *Eryngium* species. *J. Ethnopharmacol.*, **2006**, *107* (1), 32-37.
- [23] Lisciani, R.; Fattorusso, E.; Surano, V.; Cozzolino, S.; Giannatasio, M.; Sorrentino, L. Anti-inflammatory activity of *Eryngium maritimum* L. rhizome extracts in intact rats. *J. Ethnopharmacol.*, **1984**, *12* (3), 263-270.
- [24] Alkofahi, A.; Sallal, A. J.; Disi, A. M. Effect of *Eryngium creticum* on the hemolytic activities of snake and scorpion venoms. *Phytother. Res.*, **1997**, *11* (7), 540-542.
- [25] Ayoub, N. A.; Kubeczka, K. H.; Nawwar, M. A. M. An unique n-propyl sesquiterpene from *Eryngium creticum* L. (Apiaceae). *Pharmazie*, **2003**, *58* (9), 674-676.
- [26] Ebrahimzadeh, M. A.; Nabavi, S. F.; Nabavi, S. M. Antioxidant activity of leaves and inflorescence of *Eryngium caucasicum* Trautv at flowering stage. *Pharmacogn. Res.*, **2009**, *1* (6), 435-439.
- [27] Ayoub, N.; Al-Azizi, M.; Konig, W.; Kubeczka, K. H. Essential oils and a novel polyacetylene from *Eryngium yuccifolium* Michx. (Apiaceae). *Recent Prog. Med. Plants*, **2006**, *11*, 237-245.
- [28] Palá-Paúl, J.; Copeland, L. M.; Brophy, J. J.; Goldsack, R. J. Essential oil composition of *Eryngium rosulatum* P. W. Michael ined.: A new undescribed species from Eastern Australia. *Biochem. Sys. Ecol.*, **2006**, *34*, 796-801.
- [29] Palá-Paúl, J.; Pérez-Alonso, M. J.; Velasco-Negueruela, A.; Vadaré, J.; Villa, A. M.; Sanz, J.; Brophy, J. J. Essential oil composition of the different parts of *Eryngium bourgatii* Gouan from Spain. *J. Chromatogr. A*, **2005**, *1074*, 235-239.
- [30] Palá-Paúl, J.; Pérez-Alonso, M. J.; Velasco-Negueruela, A.; Vadaré, J.; Villa, A. M.; Sanz, J.; Brophy, J. J. Analysis of the essential oil composition from the different parts of *Eryngium glaciale* Boiss. from Spain. *J. Chromatogr. A*, **2005**, *1094*, 179-182.
- [31] Pino, J. A.; Rosado, A.; Fuentes, V. Chemical composition of the seed oil of *Eryngium foetidum* L. from Cuba. *J. Essential Oil Res.*, **1997**, *9*, 123-124.
- [32] Palá-Paúl, J.; Brophy, J. J.; Pérez-Alonso, M. J.; Usano, J.; Soria, S. C. Essential oil composition of the different parts of *Eryngium corniculatum* Lam. (Apiaceae) from Spain. *J. Chromatogr. A*, **2007**, *1175*, 289-293.
- [33] Chowdhury, J. U.; Nandi, N. C.; Yusuf, M. Chemical constituents of essential oil of the leaves of *Eryngium foetidum* from Bangladesh. *Bangladesh J. Sci. Ind. Res.*, **2007**, *42*, 347-352.
- [34] Thi, N.D.T.; Anh, T.H.; Thach, L.N. The essential oil composition of *Eryngium foetidum* L. in South Vietnam extracted by hydrodistillation under conventional heating and microwave irradiation. *Jeobp*, **2008**, *11*, 154-161.
- [35] Capetanos, C.; Saroglou, V.; Marin, P. D.; Simic, A.; Skaltsa, H. D. Essential oil analysis of two endemic *Eryngium* species from Serbia. *J. Serbian Chem. Soc.*, **2007**, *72*, 961-965.
- [36] Flamini, G.; Tebano, M.; Cioni, P. L. Composition of the essential oils from leafy parts of the shoots, flowers and fruits of *Eryngium amethystinum* from Amiata Mount (Tuscany, Italy). *Food Chem.*, **2008**, *107*, 671-674.
- [37] Palá-Paúl, J.; Brophy, J.J.; Goldsack, R.J.; Copeland, L.M.; Pérez-Alonso, M.J.; Velasco-Negueruela, A. Essential oil composition of the seasonal heterophyllous leaves of *Eryngium vesiculosum* from Australia. *Aust. J. Bot.*, **2003**, *51*, 497-501.
- [38] Assadian, F.; Masoudi, S.; Nematollahi, F.; Rustaiyan, A.; Larijani, K.; Mazloomifar, H. Volatile constituents of *Xanthogalum purpurascens* Ave-Lall., *Eryngium caeruleum* M.B. and *Pimpinella aurea* DC. three Umbelliferae herbs growing in Iran. *J. Essential Oil Res.*, **2005**, *17*, 243-245.
- [39] Palá-Paúl, J.; Copeland, L. M.; Brophy, J. J.; Goldsack, R. J. Essential oil composition of *Eryngium paludosum* (Moore et Betche) P. W. Michael : An endemic species from Eastern Australia. *J. Essential Oil Res.*, **2008**, *20*, 416-419.
- [40] Leclercq, P. A.; Dung, N. X.; Lo, V. N.; Toanh, N. V. Composition of the essential oil of *Eryngium foetidum* L. from Vietnam. *J. Essential Oil Res.*, **1992**, *4*, 423-424.
- [41] Cardozo, E.; Rubio, M.; Rojas, L. B.; Usubillaga, A. Composition of the essential oil from the leaves of *Eryngium foetidum* L. from the Venezuelan Andes. *J. Essential Oil Res.*, **2004**, *16*, 33-34.
- [42] Sefidkon, F.; Dabiri, M.; Alamshahi, A. Chemical composition of the essential oil of *Eryngium billardieri* F. Delaroché from Iran. *J. Essential Oil Res.*, **2004**, *16*, 42-43.
- [43] Pino, J.A.; Rosado, A.; Fuentes, V. Composition of the leaf oil or *Eryngium foetidum* L. from Cuba. *J. Essential Oil Res.*, **1997**, *9*, 467-468.
- [44] Semnani, K.M.; Azadbakht, M.; Houshmand, A. Composition of the essential oils of aerial parts of *Eryngium bungei* Boiss. and *Eryngium caeruleum* M. B. *Pharm. Sci.*, **2003**, 43-48.
- [45] Clemens, A.J.E.; Otto, S. A cyclohexenone and a cyclohexadienone glycoside from *Eryngium campestre*. *Phytochemistry*, **1986**, *25* (3), 741-743.
- [46] Muckensturm, B.; Boulanger, A.; Farahi, M.; Reduron, J.P. Secondary metabolites from *Eryngium* species. *Nat. Prod. Res.*, **2010**, *24* (5), 391-397.
- [47] Spencer, G. F. The effects of a terpene aldehyde-ester from *Eryngium paniculatum* and analogs on Velvetleaf germination. *J. Nat. Prod.*, **1986**, *49* (5), 924-926.
- [48] Brophy, J.J.; Goldsack, R.J.; Copeland, L.M.; Pala-Paul, J. Essential oil of *Eryngium* L. species from New South Wales (Australia). *J. Essential Oil Res.*, **2003**, *15*, 392-397.
- [49] Hiller, K.; Pohl, B.; Franke, P. Flavonoid spectrum of *Eryngium maritimum* L. Part 35. Components of some Saniculoideae. *Pharmazie*, **1981**, *36* (6), 451-452.
- [50] Hohmann, J.; Páll, Z.; Günther, G.; Máthé, I. Flavonolacyl glycosides of the aerial parts of *Eryngium campestre*. *Planta Med.*, **1997**, *63* (1), 96.
- [51] Ikramov, M.T.; Bandyukova, V.A.; Khalmatov, K.K. Flavonoids of some *Eryngium* species. *Him. Prir. Soedin.*, **1971**, *7* (1), 117-118.
- [52] Kartnig, T.; Wolf, J. Flavonoids from the aerial parts of *Eryngium campestre*. *Planta Med.*, **1993**, *59* (3), 285.
- [53] Nacef, S.; Ben Jannet, H.; Hamza, M. A.; Mighri, Z. Contribution to the phytochemical investigation of the plant *Eryngium dichotomum* Desf. (Apiaceae) from Tunisia. *J. de la Societe Chimique de Tunisie*, **2008**, *10* (2), 141-148.
- [54] Zarnack, J.; Hildebrandt, B.; Hiller, K.; Otto, A. To the knowledge of the compounds contained in some Saniculoideae. part XXXIII. isolation of flavonol glycosides from *Eryngium giganteum* M. B. *Zeitschrift fuer Chemie*, **1979**, *19* (6), 214-215.
- [55] Erdelmeier, C.A.; Sticher, O. Coumarin derivatives from *Eryngium campestre*. *Planta Med.*, **1985**, *51* (5), 407-409.
- [56] Mariano, P.; Mariano, P.G. Coumarins from *Eryngium ilicifolium*. *J. Nat. Prod.*, **1985**, *48* (5), 853-854.
- [57] Crowden, R.K.; Harborne, J.B.; Heywood, V.H. Chemosystematics of the umbelliferae—a general survey. *Phytochemistry*, **1969**, *8*, 1963-1984.
- [58] Pérez Gutiérrez, R.M.; Vargas Solis, R.  $\gamma$ -Lactone isolated from methanol extract of the leaves of *Eryngium carlinae* and their antispasmodic effect on rat ileum. *BLACP MA*, **2006**, *5*, 51-56.
- [59] Anam, E.M. A novel triterpenoid saponin from *Eryngium foetidum*. *Indian J. Chem. B*, **2002**, *41B* (7), 1500-1503.
- [60] Hiller, K.; Nguyen, K.Q.; Franke, P. Isolation of 3-O-D-glucopyranosyl oleanolic acid 28-O-D-xylopyranoside from *Eryngium bromeliifolium* Delar. 29. Constituents of some Saniculoideae. *Pharmazie*, **1978**, *33* (1), 78-80.
- [61] Hiller, K.; Nguyen, K.Q.C.; Franke, P. Data on the components of some Saniculoideae. XXX. Isolation of betulinic acid-3-O- $\beta$ -D-glucopyranosyl(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside from *Eryngium bromeliifolium* Delar. *Zeitschrift fuer Chemie*, **1978**, *18* (7), 260-261.
- [62] Hiller, K.; Nguyen, K.Q.C.; Franke, P.; Hintsche, R. Contents of several Saniculoideae. Part 26. Isolation of betulic acid 3-O- $\beta$ -D-glucoside, a saponin from *Eryngium bromeliifolium* Delar. *Pharmazie*, **1976**, *31* (2), 891-893.
- [63] Hiller, K.; Von Mach, B.; Franke, P. On the saponins of *Eryngium maritimum* L. Part 25. Toward information on components of some Saniculoideae. *Pharmazie*, **1976**, *31* (1), 53.
- [64] Hiller, K.; Von, T.N.; Lehmann, G.; Grundemann, E. Betulinic acid-a saponin in *Eryngium bromeliifolium* Delar. 21. The contents of a saniculoidea. *Pharmazie*, **1974**, *29* (2), 148-149.

- [65] Ikramov, M.T.; Kharlamov, I.A.; Khazanovich, R.L.; Khalmatov, K.K. Saponins of *Eryngium macrocalyx*. *Him. Prir. Soedin.*, **1976**, (3), 401.
- [66] Ikramov, M.T.; Khazanovich, R.L.; Khalmatov, K.K. Saponins from two species of *Eryngium*. *Him. Prir. Soedin.*, **1971**, 7 (6), 843.
- [67] Zhao, J.; Yang, X.W.; Hattori, M. Three new triterpene saponins from the seeds of *Aesculus chinensis*. *Chem. Pharm. Bull.*, **2001**, 49, 626-628.
- [68] Yang, X. W.; Zhao, J.; Cui, Y. X.; Liu, X. H.; Ma, C. M.; Hattori, M.; L.H., Z. Anti-HIV-1 protease triterpenoid saponins from the seeds of *Aesculus chinensis*. *J. Nat. Prod.*, **1999**, 62, 1510-1513.
- [69] Zhang, Z.Z.; Koike, K.; Jia, Z.H.; Nikaido, T.; Guo, D.A.; Zheng, J.H. New saponins from the seeds of *Aesculus chinensis*. *Chem. Pharm. Bull.*, **1999**, 47, 1515-1520.
- [70] Zhao, J.; Yang, X.W. Four new triterpene saponins from the seeds of *Aesculus chinensis*. *J. Asian Nat. Prod. Res.*, **2003**, 5, 197-203.
- [71] Wei, F.; Ma, L.Y.; Jin, W.T.; Ma, S.C.; Han, G.Z.; Khan, I.A.; Lin, R.C. Antiinflammatory triterpenoid saponins from the seeds of *Aesculus chinensis*. *Chem. Pharm. Bull.*, **2004**, 52, 1246-1248.
- [72] D'Acquarica, I.; Di Giovanni, M.C.; Gasparrini, F.; Misiti, D.; D'Arrigo, C.; Fagnano, N.; Guarnieri, D.; Iacono, G.; Bifulco, G.; Riccio, R. Isolation and structure elucidation of four new triterpenoid estersaponins from fruits of *Pittosporum tobira* AIT. *Tetrahedron*, **2002**, 58, 10127-10136.
- [73] Matsushita, A.; Miyase, T.; Noguchi, H.; Vander Velde, D. Oleanane saponins from *Sanicula elata* var. *chinensis*. *J. Nat. Prod.*, **2004**, 67, 377-383.
- [74] Voutquenne, L.; Guinot, P.; Froissard, C.; Thoison, O.; Litaudon, M.; Lavaud, C. Haemolytic acylated triterpenoid saponins from *Harpullia austro-caledonica*. *Phytochemistry*, **2005**, 66, 825-835.
- [75] Sirtori, C. R. Aescin: pharmacology, pharmacokinetics and therapeutic profile. *Pharmacol. Res.*, **2001**, 44, 183-193.
- [76] Matsuda, H.; Li, Y.; Murakami, T.; Ninomiya, K.; Yamahara, J.; Yoshikawa, M. Effects of escins Ia, Ib, IIa, and IIb from horse chestnut, the seeds of *Aesculus hippocastanum* L., on acute inflammation in animals. *Biol. Pharm. Bull.*, **1997**, 20, 1092-1095.
- [77] Zhang, Z.Z.; Li, S.Y. Cytotoxic triterpenoid saponins from the fruits of *Aesculus pavia* L. *Phytochemistry* **2007**, 68, 2075-2086.
- [78] Chan, P.K. Acylation with diangeloyl groups at C21-22 positions in triterpenoid saponins is essential for cytotoxicity towards tumor cells. *Biochem. Pharmacol.*, **2007**, 73, 341-350.
- [79] Fu, G.; Liu, Y.; Yu, S.; Huang, X.; Hu, Y.; Chen, X.; Zhang, F. Cytotoxic oxygenated triterpenoid saponins from *Symplocos chinensis*. *J. Nat. Prod.*, **2006**, 69, 1680-1686.
- [80] Zhang, Z.Z.; Li, S.Y.; Zhang, S.M.; Gorenstein, D. Triterpenoid saponins from the fruits of *Aesculus pavia*. *Phytochemistry*, **2006**, 67, 784-794.
- [81] Wang, P.; Ownby, S.; Zhang, Z.Z.; Yuan, W.; Li, S.Y. Cytotoxicity and inhibition of DNA topoisomerase I of polyhydroxylated triterpenoids and triterpenoid glycosides. *Bioorg. Med. Chem. Lett.*, **2010**, 20, 2790-2796.
- [82] Bohlmann, F.; Zdero, C. Terpene derivatives from higher plants. XII. New terpene aldehyde esters from *Eryngium* species. *Eur. J. Inorg. Chem.*, **1971**, 104 (6), 1957-1961.
- [83] Drake, D.; Lam, J. Seseli acetylene from *Eryngium bourgatii*. *Phytochemistry*, **1972**, 11 (8), 2651-2652.
- [84] Joergen, L.; Lars, P.C.; Tove, T. Acetylenes from roots of *Eryngium bourgatii*. *Phytochemistry*, **1992**, 31 (8), 2881-2882.
- [85] Assenov, I.; Gevrenova, R. D-Mannitol from *Eryngium campestre* L. *Farmatsiia (Sofia)*, **1991**, 4 (5-6), 26-28.
- [86] Ikramov, M.T.; Khazanovich, R.L.; Khalmatov, K.K. D-mannitol from *Eryngium caeruleum* and *Eryngium macrocalyx*. *Him. Prir. Soedin.*, **1969**, 5 (6), 590.
- [87] Serova, N.A. Saponin from *Eryngium incognitum*. *Meditsinskaya Promyshlennost SSSR*, **1961**, 15 (11), 26-27.
- [88] Viemont, J.D.; Goas, M. Characterization and determination of piperolic acid in some maritime dune plants. *C. R. Seances Acad. Sci. D*, **1972**, 275 (25), 2885-2887.
- [89] Abyshev, A.Z.; Denisenko, P.P.; Abyshev, D.Z.; Kerimov, Y.B. Chemical study of some species of the Caucasus flora of the umbellate family. *Farmatsiia (Moscow)*, **1977**, 26 (4), 42-44.
- [90] El-Gamal, M.H.A.; El-Bay, F.K.; El-Tawill, B.A.H.; Gadalla, K.Z. Constituents of local plants. XXI. the coumarins of *Ficus salicifolia* L. fruits, *Eryngium creticum* Lam. and *Pityranthus tortuosus* Deaf. *Egypt. J. C.*, **1978**, 18 (4), 767-772.
- [91] Suleiman, A.K. Phytochemistry of *Eryngium creticum*. *Alexandria J. Pharm. Sci.*, **1994**, 8 (1), 73-75.
- [92] Nacef, S.; Msaddek, M.; Ben Jannet, H.; Attia, S.; Chriaa, J.; Bakhrouf, A.; Mighri, Z. Isolation and structural elucidation of a steroid and a heteroside from the plant *Eryngium dichotomum*. GC/MS identification of some plants and study of their antibacterial activities. *J. Soc. Alger. Chim.*, **2003**, 13 (2), 163-175.
- [93] Emilia, K.; Ange, B.; Anna, K.; Danuta, K.; Joseph, C. New cis-chrysanthenyl esters from *Eryngium planum* L. *Nat. Prod. Commun.*, **2008**, 3 (2), 113-116.
- [94] Hiller, K.; Keipert, M.; Pfeifer, S.; Kraft, R. Components of some Saniculoideae. XI. Oligosaccharides in *Eryngium* species. *Z. Naturforsch. B*, **1969**, 24 (1), 36-38.
- [95] Hiller, K.; Keipert, M.; Pfeifer, S.; Kraft, R. Composition of some Saniculoideae. XX. Saponin spectrum of *Eryngium planum* leaves. *Pharmazie*, **1974**, 29 (1), 54-57.
- [96] Hiller, K.; Otto, A.; Gruendemann, E. Isolation of kaempferol-3-O-(6-O-β-D-glucopyranosyl)-β-D-galactopyranoside, a new flavonol glycoside from *Eryngium planum* L. Part 34: Knowledge of the constituents of some Saniculoideae. *Pharmazie*, **1980**, 35 (2), 113-114.
- [97] Leokadia, S.P. Kaempferol 3,7-dirhamnoside from *Eryngium planum* L. *Zeitschrift fuer Chemie*, **1983**, 23 (8), 294-295.
- [98] Zarnack, J.; Hiller, K.; Otto, A. The components of several Saniculoideae. XXVIII. Isolation of kaempferol 3,7-di-O-rhamnoside from *Eryngium planum* L. *Zeitschrift fuer Chemie*, **1977**, 17 (12), 445-446.
- [99] Wolfe, E.A.; Sherwood, G.A.; Mitchell, K.A.; Browne, M.P. Anti-inflammatory and analgesic properties from the stems of *Eryngium foetidum* (Shadow Benny). In *220th ACS National Meeting*, Washington, DC, United States, 2000.
- [100] Zhang, Z.Z.; Li, S.Y.; Lian, X.Y. An overview of genus *Aesculus* L.: ethnobotany, phytochemistry, and pharmacological activities. *Pharm. Crops*, **2010**, 1, 24-51.
- [101] Bogucka-Kocka, A.; Smolarz, H.D.; Kocki, J. Apoptotic activities of ethanol extracts from some Apiaceae on human leukaemia cell lines. *Fitoterapia* **2008**, 79, 487-497.
- [102] Khader, M.; Bresgen, N.; Eckl, P.M. Antimutagenic effects of ethanolic extracts from selected Palestinian medicinal plants. *J. Ethnopharmacol.*, **2010**, 127 (2), 319-324.
- [103] Strzelecka, M.; Bzowska, M.; Koziel, J.; Szuba, B.; Dubiel, O.; Riviera Nunez, D.; Heinrich, M.; Bereta, J. Anti-inflammatory effects of extracts from some traditional Mediterranean diet plants. *J. Physiol. Pharmacol.*, **2005**, 56, 1139-1156.
- [104] Jacker, H.J.; Hiller, K. The antixudative effect of saponin-5 from *Eryngium planum* L. and *Sanicular europaea* L. *Pharmazie*, **1976**, 31 (10), 747-748.
- [105] Sáenz, M.T.; Fernández, M.A.; García, M.D. Antiinflammatory and analgesic properties from leaves of *Eryngium foetidum* L. (Apiaceae). *Phytother. Res.*, **1997**, 11 (1), 380-383.
- [106] Afifi, F.U.; Al-Khalil, S.; Aqel, M.; Al-Muhteseb, M.H.; Jaghabir, M.; Saket, M.; Muheid, A. Antagonistic effect of *Eryngium creticum* extract on scorpion venom *in vitro*. *J. Ethnopharmacol.*, **1990**, 29 (1), 43-49.
- [107] Madi, J.; Munib, S.; Fatma, A.; Suleiman, A.K.; Ali, M. Potency of a Jordanian *Leiurus quinquestriatus* scorpion venom and the effect of *Eryngium creticum* on this potency. *Dirasat - Univ. Jordan*, **1989**, 16 (4), 172-178.
- [108] Ndip, R.N.; Malange Tarkang, A.E.; Mbulah, S.M.; Luma, H.N.; Agnes, M.; Ndip, L. M.; Kennedy, N.; Clare, W.; Efang, S.M.N. *In vitro* anti-helicobacter pylori activity of extracts of selected medicinal plants from North West Cameroon. *J. Ethnopharmacol.*, **2007**, 114 (3), 452-457.
- [109] Paula, J.H.A.; Seafortha, C.E.; Tikasinghb, T. *Eryngium foetidum* L.: a review. *Fitoterapia*, **2011**, 82 (3), 302-308.
- [110] Carlos, C.; Jose, G.M.; Diana, S.; Giani, S.; Felix, T.; Ange, B.; Ligia, S.; Joseph, C. Composition of a volatile extract of *Eryngium duriaei* subsp. *juresianum* (M. Lainz) M. Lainz, signalled by the antifungal activity. *Dirasat - Univ. Jordan*, **2011**, 54 (3), 619-622.
- [111] Yusuf, A.J.; Hana, S.; Abdu, S. Antimycotic activities of selected plant flora, growing wild in Lebanon, against phytopathogenic fungi. *J. Agric. Food Chem.*, **2002**, 50 (11), 3208-3213.
- [112] Laetitia, M.D.; Le Floch, G.; Magne, C. Radical scavenging, antioxidant and antimicrobial activities of *Halophytic* species. *J. Ethnopharmacol.*, **2008**, 116 (2), 258-262.

- [113] Hiller, K.; Friedrich, E.G. Antimycotic effect of Astringia-, Eryngium- and Sanicula saponins. *Pharmazie*, **1974**, *29* (12), 787-788.
- [114] Hiller, K.; Friedrich, E. Isolating antimycotically active saponin mixtures from umbellifers. *Ger. (East)*, **1975**, DD 114256 A1 19750720.
- [115] Wong, S.P.; Leong, L.P.; Koh, J.H.W. Antioxidant activities of aqueous extracts of selected plants. *Food Chem.*, **2006**, *99* (4), 775-783.
- [116] Chanwitheesuk, A.; Teerawutgulrag, A.; Rakariyatham, N. Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. *Food Chem.*, **2005**, *92* (3), 491-497.
- [117] Khaled, T.; Feras, Q. A.; Mohammad, G.; Mohammad, M.; Tamam, E. E. Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chem.*, **2007**, *104* (4), 1372-1378.
- [118] Truong Tuyet, M.; Nghiem Nguyet, T.; Pham Gia, T.; Nguyen, V. C. Alpha-glucosidase inhibitory and antioxidant activities of Vietnamese edible plants and their relationships with polyphenol contents. *J. Nutr. Sci. Vitaminol.*, **2007**, *53* (3), 267-276.
- [119] Vit, K.; Lubomir, O.; Eliska, B.; Zuzana, R.; Frantisek, C.; Katerina, K.; Kamil, K.; Daniel, J.; Miroslav, P.; Jiri, K. Evaluation of natural antioxidants of *Leuzea carthamoides* as a result of a screening study of 88 plant extracts from the European Asteraceae and Cichoriaceae. *J. Enzym. Inhib.*, **2008**, *23* (2), 218-224.
- [120] Dall'Acqua, S.; Cervellati, R.; Loi, M. C.; Innocenti, G. Evaluation of *in vitro* antioxidant properties of some traditional Sardinian medicinal plants: investigation of the high antioxidant capacity of *Rubus ulmifolius*. *Food Chem.*, **2007**, *106* (2), 745-749.
- [121] Ebrahimzadeh, M.; Nabavi, S. M.; Nabavi, S. F.; Eslami, S.; Bekhradnia, A. R. Mineral elements and antioxidant activity of three locally edible and medicinal plants in Iran. *Asian J. Chem.*, **2010**, *22* (8), 6257-6266.
- [122] Nabavi, S. M.; Ebrahimzadeh, M. A.; Nabavi, S. F.; Jafari, M. Free radical scavenging activity and antioxidant capacity of *Eryngium caucasicum* trautv and *Froripia subpinnata*. *Pharmacology*, **2008**, (3), 19-25.
- [123] Ahmet, M.; Glen, D. L.; Saban, K.; Ali, Y. Inhibition of iron-fructose-phosphate-induced lipid peroxidation in lecithin liposome and linoleic acid emulsion systems by some edible plants. *J. Food Biochem.*, **2011**, *35* (3), 833-844.
- [124] Violet, K.; Fatma, U. A.; Imad, H. Evaluation of the acute antihyperglycemic effects of four selected indigenous plants from Jordan used in traditional medicine. *Pharm. Biol.*, **2011**, *49* (7), 687-695.
- [125] Majid, S.; Katayoun, M. S. Effect of the essential oil of *Eryngium caeruleum* on percutaneous absorption of piroxicam through rat skin. *J. Essential Oil-Bearing Plants*, **2008**, *11* (5), 485-495.
- [126] Majid, S.; Katayoun, M. S. Penetration-enhancing effect of the essential oil and methanolic extract of *Eryngium bungei* on percutaneous absorption of piroxicam through rat skin. *J. Essential Oil-Bearing Plants*, **2009**, *12* (6), 728-741.
- [127] Simon, O. R.; Singh, N. Demonstration of anticonvulsant properties of an aqueous extract of Spirit Weed (*Eryngium foetidum* L.). *West Indian Med. J.*, **1986**, *35* (2), 121-125.

---

Received: December 27, 2011

Revised: February 06, 2012

Accepted: February 08, 2012

© Wang *et al.*; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.